



OPEN ACCESS

ORIGINAL ARTICLE

Genome-wide association study identifies *ERBB4* on 2q34 as a novel locus associated with sperm motility in Japanese men

Youichi Sato,¹ Atsushi Tajima,^{2,3} Takehiro Sato,³ Shiari Nozawa,⁴ Miki Yoshiike,⁴ Issei Imoto,² Aiko Yamauchi,¹ Teruaki Iwamoto^{4,5}

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/jmedgenet-2017-104991>).

¹Department of Pharmaceutical Information Science, Graduate School of Biomedical Sciences, Tokushima University, Tokushima, Japan

²Department of Human Genetics, Graduate School of Biomedical Sciences, Tokushima University, Tokushima, Japan

³Department of Bioinformatics and Genomics, Graduate School of Advanced Preventive Medical Sciences, Kanazawa University, Kanazawa, Japan

⁴Department of Urology, St. Marianna University School of Medicine, Kawasaki, Japan

⁵Center for Infertility and IVF, International University of Health and Welfare Hospital, Nasushiobara, Japan

Correspondence to

Dr Youichi Sato, Department of Pharmaceutical Information Science, Graduate School of Biomedical Sciences, Tokushima University, 1-78-1 Sho-machi, Tokushima, 770-8505, Japan; youichi.sato@tokushima-u.ac.jp and Prof. Atsushi Tajima, Department of Bioinformatics and Genomics, Graduate School of Advanced Preventive Medical Sciences, Kanazawa University, 13-1 Takara-machi, Kanazawa, 920-8640, Japan; atajima@med.kanazawa-u.ac.jp

YS and AT contributed equally.

Received 20 August 2017
Revised 29 December 2017
Accepted 21 January 2018
Published Online First
16 February 2018



To cite: Sato Y, Tajima A, Sato T, et al. *J Med Genet* 2018;**55**:415–421.

ABSTRACT

Background The decrease in sperm motility has a potent influence on fertilisation. Sperm motility, represented as the percentage of motile sperm in ejaculated sperms, is influenced by lifestyle habits or environmental factors and by inherited factors. However, genetic factors contributing to individual differences in sperm motility remain unclear. To identify genetic factors that influence human sperm motility, we performed a genome-wide association study (GWAS) of sperm motility.

Methods A two-stage GWAS was conducted using 811 Japanese men in a discovery stage, followed by a replication study using an additional 779 Japanese men.

Results In the two-staged GWAS, a single nucleotide polymorphism rs3791686 in the intron of gene for erb-b2 receptor tyrosine kinase 4 (*ERBB4*) on chromosome 2q34 was identified as a novel locus for sperm motility, as evident from the discovery and replication results using meta-analysis ($\beta=-4.01$, combined $P=5.40\times 10^{-9}$).

Conclusions Together with the previous evidence that Sertoli cell-specific *ErbB4*-knockout mice display an impaired ability to produce motile sperm, this finding provides the first genetic evidence for further investigation of the genome-wide significant association at the *ERBB4* locus in larger studies across diverse human populations.

INTRODUCTION

Approximately 10% couples display infertility issues, and half of these problems are related to men.^{1,2} Male factor infertility may arise from various medical conditions such as spermatogenic failure, varicocele, obstructive azoospermia and congenital absence of vas deferens. Sperm motility—represented as the percentage of motile sperm in the ejaculated sperms—has a large influence on the fertilisation ability. Therefore, several studies are conducted to understand the factors that affect sperm motility.

Oxidative stress induced by alcohol consumption, cigarette smoking, obesity, diabetes, physical exercise, psychological stress, ageing, infection and environment factors (pollutants such as nitric oxide, lead and electromagnetic waves from cell phones) is one of the major factors responsible for the reduction in sperm motility.^{3,4} Genetic background has also been shown to be associated

with sperm motility. The *gr/gr* subdeletion in the azoospermia factor c region of the Y chromosome was shown to be strongly associated with decreased sperm motility in men from Japanese population.⁵ Furthermore, polymorphisms in genes encoding cytochrome P450 family 19 subfamily A polypeptide 1,⁶ androgen receptor,⁷ follicle-stimulating hormone receptor,⁸ steroid 5 α -reductase⁹ and oestrogen receptor^{10,11} were associated with sperm motility. These genes are related to the reproductive hormones and contribute to the testicular development and spermatogenesis; these genes have been proposed based on their functions. However, the genetic determinants for human sperm motility are poorly understood.

Genome-wide association study (GWAS) is an approach to find the genetic variations associated with disease or quantitative traits. To date, four GWASs associated with male infertility have been reported. These include the non-obstructive azoospermia or oligozoospermia in Caucasians or Chinese men^{12–15} and the family size or birth rate in Hutterite men in the USA.¹⁶ In the latter, 9 of the 41 single nucleotide polymorphisms (SNPs) were significantly correlated with the family size or birth rate and found to be associated with reduced sperm quantity and/or function in the subsequent validation study using 123 ethnically diverse men. However, there are no reports on GWAS of sperm motility. Here, we clarified the genetic determinants for human sperm quality by conducting a GWAS of sperm motility in 811 Japanese men, with a subsequent validation of the association in an additional 779 Japanese men.

METHODS

Subjects

We performed a two-staged genetic association study. The discovery stage included 816 men (20.7 \pm 1.7 years old, mean \pm SD) from the young Japanese population. These were recruited from university students in three study centres based in departments of urology at university hospitals in Japan (Kawasaki, Kanazawa and Nagasaki) as previously reported.¹⁷ The inclusion criteria were that the man was 18–24 years and that both he and his mother were born in Japan. The replication stage included 779 men (31.2 \pm 4.8 years old, mean \pm SD) of proven fertility recruited from the partners of pregnant women who attended obstetric clinics in

Table 1 Characteristics of subjects

	Discovery (n=811)	Replication (n=779)	P value
Age (years)	20.7±1.7	31.2±4.8	<0.0001
BMI (kg/m ²)	21.5±2.5	23.3±3.0	<0.0001
Ejaculation abstinence (hours)	79.8±39.5	194.6±326.8	<0.0001
Sperm motility (%)	60.3±14.5	59.4±20.7	0.32

Data are presented as mean±SD. P values were obtained with Student's unpaired t-test.

BMI, body mass index.

four cities in Japan (Sapporo, Kanazawa, Osaka and Fukuoka).¹⁸ The inclusion criteria for the men were as follows: age 20–45 years and both he and his mother had to be born and live in Japan. In addition, the current pregnancy of the female partner had to be achieved by normal sexual relations and not as a result of fertility treatment. We excluded sample with complete deletion of AZF region in both subjects. The characteristics of the two-staged subjects are summarised in table 1 Table 1. No difference in sperm motility was observed between the two-staged subjects. Some of the subjects in this study have been described in previous reports.^{19–26}

Clinical trait measurements

The measurement of clinical trait of these subjects has been described in previous reports.^{17, 18} Briefly, age, body weight, height and ejaculation abstinence period were self-reported. Body mass index (BMI) (kg/m²) was calculated from the body weight and height. Semen samples were obtained once by masturbation after sexual abstinence for at least 48 hours and ejaculated into clean, wide-necked, sterile, non-toxic collection containers. The samples were protected from extremes of temperature and liquefied at 37°C prior to their examination. At each semen collection site, sperm motility was assessed from 10 µL of well-mixed semen, which was placed on a clean glass slide, covered and examined at a total magnification of 400× at 37°C. Sperm motility (%) was calculated as ([number of motile sperm in the ejaculate]/[number of sperm in the ejaculate])×100. The motility assessment was repeated on a second 10 µL aliquot of semen and the average value calculated. Sperms were assessed using the WHO motility classes A, B, C and D,²⁷ wherein sperms from classes A and B were considered as motile. Technicians from each centre were initially trained by one technician from St. Marianna University in Kawasaki, and these clinical trait measurements were similarly performed in both cohorts.

Genotyping, quality control and imputation

Genomic DNA was extracted from the peripheral blood samples of subjects using a QIAamp DNA blood kit (Qiagen, Tokyo, Japan). In the discovery stage, 816 men were genotyped using the Illumina HumanCore V.1.0 DNA Analysis Kit (Illumina, Tokyo, Japan) following the manufacturer's instructions. We genotyped 298 930 SNPs, and the quality control of genotyped SNPs and samples was conducted using PLINK V.1.07 software package (<http://pngu.mgh.harvard.edu/~purcell/plink/>).²⁸ Of the 816 samples, four were excluded because these were duplicates or familial relationships (PI_HAT>0.25), as revealed by pairwise identical-by-state/identity-by-descent estimation. Furthermore, we excluded one sample that was identified as a genetic outlier by the principle component analysis-based method using the genotype data of the HapMap CHB and JPT as the internal

controls (online supplementary figure S1). Finally, 811 samples were included for genome-wide association analysis.

For genotype imputation analysis, only non-redundant polymorphic SNPs with reference SNP (rs) IDs fulfilling the following criteria were included: (1) per-SNP call rate ≥0.98 and (2) P value for Hardy-Weinberg equilibrium (HWE) ≥10⁻⁶ in our sample set. Genotype data were flipped to forward strand with conform-gt, which is the utility program of BEAGLE V.4.1,^{29, 30} using genotype data for Asian samples (JPT and CHB) of the 1000 Genomes Project^{31, 32} as a reference panel. Imputation was performed with BEAGLE V.4.1, using the 1000 Genomes Project Phase 3 V.5 as a reference panel. We excluded SNPs with R² <0.8 and all indels from the imputed genotype data to obtain genotypes for 3 901 256 SNPs, which were used for subsequent association analyses.

In the replication stage, SNP rs3791686 was genotyped using TaqMan probe (C_27517144_10; Applied Biosystems, Tokyo, Japan) with the ABI 7900HT real-time PCR system (Applied Biosystems). rs3791686 in randomly selected 100 samples of discovery subject was directly genotyped to confirm the concordance of the imputed results. The concordance of typing results between genotyped and imputed was 100%. The genotypes of rs3791686 were in HWE in a total of 1590 samples.

Statistical analysis

In discovery and replication stages, associations between each SNP and sperm motility were assessed using a multiple linear regression under an additive genetic model, with adjustments for age, BMI, ejaculation abstinence period and time from masturbation to semen evaluation using PLINK or R V.3.1.2 software package (<http://www.R-project.org/>). Since the raw value was closest to the normal distribution than some converted values, we decided to use the raw value for analysis in the present study. We set a suggestive threshold of P values <1×10⁻⁶ in the discovery stage. The results were combined in a meta-analysis using the meta package for the R software. The extent of heterogeneity among studies was quantified by the I² statistic³³ and statistically assessed by the Cochran's Q test. No heterogeneity was observed in this study, as determined by the I² statistic <50% or P value >0.1; hence, a fixed-effect model using the inverse variance method was used. Genome-wide statistical significance was considered at P values <5×10⁻⁸.

The Manhattan and quantile–quantile plots were generated using qqman package for the R software, while a regional plot was created by LocusZoom using the 1000 Genomes project Asian (ASN) data (November 2014).³⁴ With the exception of annotations, linkage disequilibrium (LD) was calculated using PLINK software V.1.07 with genotype imputation data. Significant expression quantitative trait loci (eQTL) by SNP was searched on GTEx Portal database (<http://www.gtexportal.org/home/>).³⁵ HaploReg V.4.1 (<http://archive.broadinstitute.org/mammals/haploreg/haploreg.php>) was used for functional annotation analysis of variants,³⁶ and RegulomeDB (<http://regulome.stanford.edu/index>) was employed to identify potential regulatory functions.³⁷ Pathway analysis of GWAS datasets for sperm motility was conducted using iGSEA4GWAS V.2 (<http://gesea4gwas-v2.psych.ac.cn/>).³⁸ The SNPs were mapped to the nearest genes within 20 kb upstream/downstream and searched KEGG (<http://www.genome.jp/kegg/>), BioCarta (<http://www.biocarta.com>) and GO (<http://www.geneontology.org>) gene set/pathway databases. The threshold of false discovery rate (FDR)<0.05 is statistically significant in this analysis.




Figure 1 Manhattan plot of associations from the discovery stage of GWAS of sperm motility. The negative \log_{10} -transformed P values (Y axis) of genotyped and imputed SNPs are shown according to their positions on chromosome. The horizontal line represents suggestive (blue) and genome-wide (red) significance thresholds. GWAS, genome-wide association study; SNPs, single nucleotide polymorphisms.

RESULTS

We conducted a two-staged GWAS to identify genetic loci associated with human sperm motility. We enrolled 816 Japanese men from the university students for the discovery stage and 779 Japanese men from the partners of pregnant women for the replication stage of GWAS. After quality control of samples using initially genotyped 298 930 SNP data in the discovery stage, 811 Japanese men were selected. We performed imputation analysis, which provided typed and imputed genotypes for 3 901 256 SNPs that passed quality control. Finally, 811 samples and 3 901 256 SNPs were included for the discovery stage. The characteristics of subjects are presented in [table 1](#).

We performed GWAS between a total of 3 901 256 SNPs and sperm motility in 811 men in the discovery stage. Manhattan and quantile–quantile plots of GWAS are presented in [figure 1](#) and online supplementary figure S2, respectively. The genomic inflation factor (λ) was reported to be 1.0, indicating the unlikelihood of the inflation of the false-positive association. The top 50 GWAS candidate SNPs for sperm motility were presented in online supplementary table S1. We failed to find any SNPs to reach a genome-wide significance level ($P < 5 \times 10^{-8}$) in the discovery stage. When setting a suggestive significance threshold of P values $< 1 \times 10^{-6}$, we identified that two SNPs, rs3791686 and rs1836719 on 2q34, were suggestively associated with sperm motility ($\beta = -4.25$, discovery $P = 4.47 \times 10^{-7}$; $\beta = -4.22$, discovery $P = 5.29 \times 10^{-7}$, respectively) (online supplementary table S1). These two SNPs are in strong LD ($r^2 = 0.99$); thus, we selected only the most significant SNP (rs3791686) for the subsequent replication genotyping.

In the replication study involving 779 proven fertile men, SNP rs3791686 on 2q34 showed a significant association with sperm motility ($\beta = -3.51$, replication $P = 3.88 \times 10^{-3}$) ([table 2](#)). When we combined the discovery and replication results using meta-analysis, rs3791686 surpassed the threshold

for genome-wide significance ($\beta = -4.01$, combined $P = 5.40 \times 10^{-9}$), with no evidence of heterogeneity between the two studies. The variance in sperm motility explained by rs3791686 was 2.0%.

[Figure 2](#) shows a regional association plot for the genomic region 400 kb upstream and downstream of the lead SNP rs3791686 in the discovery stage. Within the region, 24 genotyped and 289 imputed SNPs, including rs3791686, were associated with sperm motility, with discovery P values < 0.05 from the association analysis in discovery stage (online supplementary table S2). The sperm motility-associated genomic interval indexed by rs3791686 on 2q34 overlapped with a single known gene, erb-b2 receptor tyrosine kinase 4 (*ERBB4*), while the lead SNP rs3791686 was located in the intron of *ERBB4*. Of a total of 313 SNPs with discovery P values < 0.05 within the associated interval, none resulted in amino acid substitution or protein truncation or affected the splicing of *ERBB4*; one synonymous SNP (rs3748962) and seven SNPs in the 3'-untranslated region of *ERBB4* were observed (online supplementary table S2). To obtain putative functional annotations of rs3791686 and other 13 SNPs in high LD ($r^2 > 0.80$ in East Asians from the 1000 Genomes Project) with rs3791686 (online supplementary table S3) within the associated interval, we used the following three databases: GTEx Portal,³⁵ HaploReg³⁶ and RegulomeDB.³⁷ We assessed if the 14 SNPs, including rs3791686 on 2q34, were involved in eQTLs using the GTEx Portal database and found that no significant eQTLs were associated with all the SNPs examined. HaploReg and RegulomeDB databases search revealed that the 14 SNPs examined within the associated interval may be regarded as candidate regulatory SNPs (online supplementary table S3). In the HaploReg database, the lead SNP rs3791686 itself was associated with enhancer histone marks and DNase I hypersensitive region in embryonic stem-derived cells and resided in regulatory motifs of four transcription factors—Maf, Nkx2, Nkx3 and TATA-binding protein (online supplementary table S3). Of the 13 SNPs in high LD with the lead SNP rs3791686, five were associated with enhancer histone marks and/or DNase I hypersensitive regions in various types of cells and tissues, while 12 SNPs had the potential to alter nucleotide sequences of several regulatory motifs. The RegulomeDB database provided the experimental evidence that three SNPs (rs13003941, rs1836720 and rs1836719) were located in DNase I hypersensitive and/or TF-binding regions in various cells. The iGSEA4GWAS analysis identified 421 significant pathway (FDR < 0.05) (online supplementary table S4). Numerous pathways were identified in this analysis; this finding suggests that sperm motile ability is likely to affect by a complicated process involving interaction between multiple genes and pathways.

DISCUSSION

In the first two-staged GWAS of sperm motility in Japanese men, we identified a novel sperm motility-associated locus at

Table 2 Genome-wide significant SNP identified in GWAS for sperm motility

SNP (effect/reference)	Chr.	Position (hg 19)	Gene locus	Stage	Genotype	EAF	Beta (SE)	P value	Var (%)*
rs3791686 (C/T)	2	212 221 870	<i>ERBB4</i> intron	Discovery	40/307/464	0.24	-4.25 (0.84)	4.47×10^{-7}	
				Replication	48/268/463	0.23	-3.51 (1.21)	3.88×10^{-3}	
				Meta-analysis			-4.01 (0.69)	5.40×10^{-9}	2.0

Data are shown as the estimated linear regression statistic beta, SE and P value using an additive genetic model with adjustments for age, BMI, ejaculation abstinence and time from masturbation to test.

*Var (%): percentage of phenotypic variance explained by SNP.

BMI, body mass index; Chr, chromosome; EAF, effect allele frequency; GWAS, genome-wide association study; SNP, single nucleotide polymorphism.

ERBB4 on chromosome 2q34. The most strongly associated SNP was typed by imputation analysis. In this study, the subjects of discovery stage were genotyped using the Illumina HumanCore V1.0 DNA Analysis Kit with a total of 298 930 SNPs. Subsequently, to enhance the coverage, untyped SNPs were imputed. Sometimes, imputation methods may be less accurate for typing of SNPs. To confirm the accuracy of this imputation method, randomly selected samples were directly genotyped for the GWAS-lead SNP rs3791686. The result of imputation analysis was validated by the genotyping.

SNP rs3791686 lies in the intron of *ERBB4* gene, which is a member of the receptor tyrosine kinase family and epidermal growth factor receptor subfamily. *ERBB4* is expressed in several tissues, including kidney, breast, cerebrum, heart, bone, ovary and testis. On activation by its ligands, *ERBB4* forms a dimer on the cell surface. Following cleavage of the *ERBB4* ectodomain by a disintegrin and metalloprotease domain 17 (ADAM17) and γ -secretase, the intracellular domain of *ERBB4* is translocated into the nucleus. Inside the nucleus, *ERBB4* is involved in the regulation of cell proliferation and differentiation.^{39–42} *ERBB4* is thought to be both necessary and sufficient to trigger an antiproliferative response in human breast cancer cells.⁴³ Kim *et al*⁴⁴ reported that the SNP rs13393577 in *ERBB4* is associated with breast cancer risk in Koreans by GWAS. In addition, previous GWASs in the National Human Genome Research Institute (NHGRI) GWAS Catalog demonstrate that SNPs in *ERBB4* are genome-wide significantly associated with polycystic ovary syndrome (lead SNP rs1351592)⁴⁵ and BMI (lead SNP rs7599312).⁴⁶ The lead SNPs at *ERBB4* from the previous GWASs are >1 Mb distally localised from the sperm motility-lead SNP rs3791686 and show no pairwise LD ($r^2 < 0.01$ in

East Asians) with rs3791686. This indicates a novel association for sperm motility at *ERBB4* on 2q34, which is independent of other human diseases and traits.

The expression of *ERBB4* is evident in male reproductive tissues, including testis. In the testicular tissue, *ERBB4* is expressed in both somatic cells (Sertoli cells and Leydig cells) and germ cells.⁴⁷ It is notable that Sertoli cell-specific *ErbB4*-knockout mice exhibit a developmental defect in the organisation of the testicular seminiferous tubules, which reduces male fertility. Aberration in the testicular cell adhesion machinery caused by *ErbB4* deficiency leads to a compromised capacity of the testes to produce motile sperms.⁴⁷ Thus, *ERBB4* signalling in the Sertoli cells may influence the sperm motility, suggestive of the promising functional role of *ERBB4* in sperm motility. The lead SNP rs3791686 identified in this GWAS is an intronic SNP of *ERBB4* and displays the potential to act as a functional regulatory SNP based on the multiple functional annotations. As the functional annotation analyses reveal an association between other SNPs in high LD with rs3791686 and potential regulatory domains and motifs, the sperm motility locus at *ERBB4* may have a role in the regulation of *ERBB4* expression via a cis-regulatory mechanism. Sandholm *et al*,⁴⁸ reported that a cis-eQTL for *ERBB4* in tubulointerstitial-enriched kidney biopsies maps to intronic *ERBB4* SNPs, rs17418640 and rs17418814. Both of these SNPs are proxies for rs7588550, representing a suggestive association with diabetic nephropathy; however, these eQTL SNPs are not in LD ($r^2 < 0.01$ in East Asians) with the sperm motility-lead SNP rs3791686. Further studies are warranted to assess the potential contribution of the sperm motility-associated locus indexed by rs3791686 to the regulation of *ERBB4* expression. These studies will also help explore the possible involvement of this locus in




Figure 2 Regional association plot for a sperm motility-associated locus on chromosome 2q34. The negative \log_{10} -transformed P values (Y axis) of genotyped and imputed SNPs that are located in 400 kb upstream and downstream regions of the GWAS-lead SNP rs3791686 in the discovery stage are shown according to their chromosomal positions. Purple diamond and circles represent the lead SNP rs3791686 and other SNPs within the region, respectively, with the colour of each circle indicating the range of pairwise r^2 value with rs3791686. The right Y axis shows the recombination rates estimated from the 1000 Genomes project Asian (ASN) data (November 2014). The RefSeq gene, *ERBB4*, within the region is shown in the panel below. SNPs, single nucleotide polymorphisms.

the expression regulation on a genome-wide scale via transregulatory mechanisms.

Liu *et al.*⁴⁹ have reported that five SNPs (rs215702, rs6476866, rs10129954, rs2477686 and rs10841496) were significantly correlated with sperm progressive motility. However, present study did not detect the variants associated with sperm motility including the region 400 kb upstream and downstream of these five SNPs. Previously, we also have reported that four SNPs as being significantly associated with risk factors for non-obstructive azoospermia (NOA) by Chinese GWAS¹³ were not associated with NOA in Japanese population.²¹ The reason for these may be that there are small genetic differences between Han Chinese and Japanese population by a principal component analysis using genotype data of the HapMap CHB and JPT (online supplementary figure S1). Additionally, we found a strong association between Y-haplogroup and sperm motility in the same Japanese populations.²² However, none of the SNPs on Y chromosome display a significant association ($P < 0.05$) with sperm motility in this study. The Illumina Human Core V.1.0 DNA analysis kit includes 1943 Y-chromosome markers. However, of these, only 177 markers could be examined in the discovery stage. Because this kit does not include Japanese Y-haplogroup specific markers, we did not find a significant association between Y-chromosome variants and sperm motility in this study.

Several limitations of this study should be noted. In this study, men of proven fertility were used, instead of randomly selected subjects as the replication samples. These were the only samples available for the current replication analysis. Using samples selected on the basis of fertility may cause bias. In fact, abstinence periods were significantly different between two cohorts (table 1). In general, longer abstinence period is correlated with lower sperm motility. As the previous study described, abstinence period was negatively correlated with sperm motility in both cohorts.²² To reduce the influence of the abstinence period on sperm motility, we included this as a covariate for a multiple linear regression analysis. Therefore, we think that the effect of abstinence period on the power to detect sperm motility-associated SNPs is minimized in this study. Additionally, all the participants in the current two-staged GWAS were Japanese men. Independent validation studies are required to test the observed association between *ERBB4* SNPs and sperm motility using other general populations and ethnicities. The transethnic association analyses at the *ERBB4* locus will also enable us to narrow the association signal to smaller sets of SNPs, when leveraging differences in LD structures across diverse populations. The limited statistical power of this two-staged GWAS prevented the detection of other true positive associations at a genome-wide significance level because the sample size was not large. We believe that other genetic loci may account for the interindividual variation in sperm motility, and therefore, larger scale GWAS analyses may be expected to identify novel associations between genetic variants and sperm motility.

It is one of the limitations that sperm motility may show sometimes intraindividual variation between samples from the same individual. When phenotypic repeatability is low, setting the upper boundary of heritability of a trait may decrease sensitivity to detect genetic variant/variants associated with a trait. As aforementioned, sperm motility depends on the abstinence period; in general, abstinence period and sperm motility shows a negative relationship. In our samples, although there is a difference in the strength of association, the abstinence period was indicated to be negatively correlated with sperm motility in both cohorts,²² which is not contradictory. In this study, we set a significance threshold of P values $< 1 \times 10^{-6}$ in discovery

stage and performed the replication analysis of the selected SNP. The strength of the SNP-trait association between cohorts was slightly different, but there was no significant heterogeneity. As well as intraindividual variation of sperm motility between individual samples, the measurement of sperm motility may have variability by operators (individual technicians). To reduce the between-centre variability, technicians from each centre were initially trained by one technician from St. Marianna University in Kawasaki. In addition, to statistically reduce the influence of differences in sperm assessment between the centres, we added each centre as a covariate and further conducted an association analysis between sperm motility and rs3791686. We found that rs3791686 was associated with sperm motility in the discovery stage ($\beta = -4.35$, $P = 1.62 \times 10^{-7}$) and in the replication stage ($\beta = -3.16$, $P = 0.012$). When we combined two results using meta-analysis, rs3791686 was genome-wide significantly associated with sperm motility ($\beta = -3.99$, $P = 6.60 \times 10^{-9}$). This finding was very similar to the result (table 2) from the association analysis without adjustment for semen analysis centre. Although the measurements of the semen analysis may not necessarily be representatives of individual sperm motility, together with the previous finding of Sertoli cell-specific *ErbB4*-knockout mice, we are confident that the results of our GWAS are valid.

In conclusion, this first two-staged GWAS for sperm motility identifies a novel sperm motility-associated locus at *ERBB4* on 2q34. The genetic evidence suggests that *ERBB4* is a promising candidate for future association studies in diverse populations with larger sample sizes. Further studies such as fine-scale genetic mapping are needed to uncover a functional variant at this locus as well as the underlying molecular mechanism.

Acknowledgements We thank all the volunteers who participated in this study. We are grateful to the late Professor Yutaka Nakahori and Professors Eitetsue Koh, Jiro Kanaya, Mikio Namiki, Kiyomi Matsumiya, Akira Tsujimura, Kiyoshi Komatsu, Naoki Itoh and Jiro Eguchi for collecting blood samples from the participants. We also thank Professor Toyomasa Katagiri for his assistance with the AB GeneAmp PCR System 9700.

Contributors YS and AT conceived, designed the experiments, performed the experiments and wrote the paper. TS performed the imputation analysis. SN, MY and TI prepared and collected samples. II contributed material and analysis tools. YS, AT, TS, II, AY and TI reviewed and revised the manuscript.

Funding This study was supported in part by the Ministry of Health and Welfare of Japan (1013201) (to TI), Grant-in-Aids for Scientific Research (C) (26462461) (to YS), (23510242) (to AT) and Grant-in-Aids for Scientific Research (B) (17H04331) (to YS), (15H04320) (to AT) from the Japan Society for the Promotion of Science, the European Union (BMH4-CT96-0314) (to TI), the Takeda Science Foundation (to AT) and The Suzuki Urinary Foundation (to YS).

Competing interests None declared.

Patient consent Obtained.

Ethics approval This study was approved by the ethics committees of the University of Tokushima and St. Marianna Medical University. All participants provided written informed consent.

Provenance and peer review Not commissioned; externally peer reviewed.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2018. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- Skakkebaek NE, Giwercman A, de Kretser D. Pathogenesis and management of male infertility. *Lancet* 1994;343:1473–9.

- 2 McLachlan RI, de Kretser DM. Male infertility: the case for continued research. *Med J Aust* 2001;174:116–7.
- 3 Agarwal A, Virk G, Ong C, du Plessis SS. Effect of oxidative stress on male reproduction. *World J Mens Health* 2014;32:1–17.
- 4 Opuwari CS, Henkel RR. An update on oxidative damage to spermatozoa and oocytes. *Biomed Res Int* 2016;2016:1–11.
- 5 Sato Y, Iwamoto T, Shinka T, Nozawa S, Yoshiike M, Koh E, Kanaya J, Namiki M, Matsumiya K, Tsujimura A, Komatsu K, Itoh N, Eguchi J, Yamauchi A, Nakahori Y. Y Chromosome gr/gr subdeletion is associated with lower semen quality in young men from the general Japanese population but not in fertile Japanese men. *Biol Reprod* 2014;90:116.
- 6 Lazaros L, Xita N, Kaponis A, Hatz E, Plachouras N, Sofikitis N, Zikopoulos K, Georgiou I. The association of aromatase (CYP19) gene variants with sperm concentration and motility. *Asian J Androl* 2011;13:292–7.
- 7 Delli Muti N, Agarwal A, Buldregghini E, Gioia A, Lenzi A, Boscaro M, Balercia G. Have androgen receptor gene CAG and GGC repeat polymorphisms an effect on sperm motility in infertile men? *Andrologia* 2014;46:564–9.
- 8 Lazaros L, Xita N, Takenaka A, Sofikitis N, Makrydimas G, Stefanos T, Kosmas I, Zikopoulos K, Hatz E, Georgiou I. Synergistic effect of follicle-stimulating hormone receptor and androgen receptor gene variants on semen quality. *Andrologia* 2013;45:339–44.
- 9 Zhao D, Wu W, Xu B, Niu X, Cui H, Zhang Y, Wang Z, Wang X. Variants in the SRD5A2 gene are associated with quality of semen. *Mol Med Rep* 2012;6:639–44.
- 10 Lazaros LA, Xita NV, Kaponis AI, Zikopoulos KA, Plachouras NI, Georgiou IA. Estrogen receptor alpha and beta polymorphisms are associated with semen quality. *J Androl* 2010;31:291–8.
- 11 Safarinejad MR, Shafiei N, Safarinejad S. Association of polymorphisms in the estrogen receptors alpha, and beta (ESR1, ESR2) with the occurrence of male infertility and semen parameters. *J Steroid Biochem Mol Biol* 2010;122:193–203.
- 12 Aston KI, Carrell DT. Genome-wide study of single-nucleotide polymorphisms associated with azoospermia and severe oligozoospermia. *J Androl* 2009;30:711–25.
- 13 Hu Z, Xia Y, Guo X, Dai J, Li H, Hu H, Jiang Y, Lu F, Wu Y, Yang X, Li H, Yao B, Lu C, Xiong C, Li Z, Gui Y, Liu J, Zhou Z, Shen H, Wang X, Sha J. A genome-wide association study in Chinese men identifies three risk loci for non-obstructive azoospermia. *Nat Genet* 2012;44:183–6.
- 14 Zhao H, Xu J, Zhang H, Sun J, Sun Y, Wang Z, Liu J, Ding Q, Lu S, Shi R, You L, Qin Y, Zhao X, Lin X, Li X, Feng J, Wang L, Trent JM, Xu C, Gao Y, Zhang B, Gao X, Hu J, Chen H, Li G, Zhao J, Zou S, Jiang H, Hao C, Zhao Y, Ma J, Zheng SL, Chen ZJ. A genome-wide association study reveals that variants within the HLA region are associated with risk for nonobstructive azoospermia. *Am J Hum Genet* 2012;90:900–6.
- 15 Ni B, Lin Y, Sun L, Zhu M, Li Z, Wang H, Yu J, Guo X, Zuo X, Dong J, Xia Y, Wen Y, Wu H, Li H, Zhu Y, Ping P, Chen X, Dai J, Jiang Y, Xu P, Du Q, Yao B, Weng N, Lu H, Wang Z, Zhu X, Yang X, Xiong C, Ma H, Jin G, Xu J, Wang X, Zhou Z, Liu J, Zhang X, Conrad DF, Hu Z, Sha J. Low-frequency germline variants across 6p22.2–6p21.33 are associated with non-obstructive azoospermia in Han Chinese men. *Hum Mol Genet* 2015;24:5628–36.
- 16 Kosova G, Scott NM, Niederberger C, Prins GS, Ober C. Genome-wide association study identifies candidate genes for male fertility traits in humans. *Am J Hum Genet* 2012;90:950–61.
- 17 Iwamoto T, Nozawa S, Mieno MN, Yamakawa K, Baba K, Yoshiike M, Namiki M, Koh E, Kanaya J, Okuyama A, Matsumiya K, Tsujimura A, Kanetake H, Eguchi J, Skakkebaek NE, Vierula M, Toppari J, Jørgensen N. Semen quality of 1559 young men from four cities in Japan: a cross-sectional population-based study. *BMJ Open* 2013;3:e002222.
- 18 Iwamoto T, Nozawa S, Yoshiike M, Namiki M, Koh E, Kanaya J, Okuyama A, Matsumiya K, Tsujimura A, Komatsu K, Tsukamoto T, Itoh N, Mieno MN, Vierula M, Toppari J, Skakkebaek NE, Jørgensen N. Semen quality of fertile Japanese men: a cross-sectional population-based study of 792 men. *BMJ Open* 2013;3:e002223.
- 19 Nakahori Y, Sato Y, Ewis AA, Iwamoto T, Shinka T, Nozawa S, Yoshiike M, Yang XJ, Sei M, Namiki M, Kou E, Ito N, Komatsu K, Matsumiya K, Nakagome Y. Climatic influence on the reproductive characteristics of Japanese males. *J Hum Genet* 2012;57:375–8.
- 20 Sato Y, Shinka T, Iwamoto T, Yamauchi A, Nakahori Y. Y chromosome haplogroup D2* lineage is associated with azoospermia in Japanese males. *Biol Reprod* 2013;88:107.
- 21 Sato Y, Jinam T, Iwamoto T, Yamauchi A, Imoto I, Inoue I, Tajima A. Replication study and meta-analysis of human non-obstructive azoospermia in Japanese populations. *Biol Reprod* 2013;88:87.
- 22 Sato Y, Iwamoto T, Shinka T, Nozawa S, Yoshiike M, Koh E, Kanaya J, Namiki M, Matsumiya K, Tsujimura A, Komatsu K, Itoh N, Eguchi J, Yamauchi A, Nakahori Y. Y chromosome gr/gr subdeletion is associated with lower semen quality in young men from the general Japanese population but not in fertile Japanese men. *Biol Reprod* 2014;90:116.
- 23 Sato Y, Shinka T, Nozawa S, Yoshiike M, Koh E, Kanaya J, Namiki M, Matsumiya K, Tsujimura A, Komatsu K, Itoh N, Eguchi J, Yamauchi A, Iwamoto T, Nakahori Y. Y chromosome haplogroup D2a1 is significantly associated with high levels of luteinizing hormone in Japanese men. *Andrology* 2015;3:520–5.
- 24 Sato Y, Tajima A, Tsunematsu K, Nozawa S, Yoshiike M, Koh E, Kanaya J, Namiki M, Matsumiya K, Tsujimura A, Komatsu K, Itoh N, Eguchi J, Imoto I, Yamauchi A, Iwamoto T. Lack of replication of four candidate SNPs implicated in human male fertility traits: a large-scale population-based study. *Hum Reprod* 2015;30:1505–9.
- 25 Sato Y, Tajima A, Tsunematsu K, Nozawa S, Yoshiike M, Koh E, Kanaya J, Namiki M, Matsumiya K, Tsujimura A, Komatsu K, Itoh N, Eguchi J, Imoto I, Yamauchi A, Iwamoto T. An association study of four candidate loci for human male fertility traits with male infertility. *Hum Reprod* 2015;30:1510–4.
- 26 Sato Y, Tajima A, Katsurayama M, Nozawa S, Yoshiike M, Koh E, Kanaya J, Namiki M, Matsumiya K, Tsujimura A, Komatsu K, Itoh N, Eguchi J, Imoto I, Yamauchi A, Iwamoto T. A replication study of a candidate locus for follicle-stimulating hormone levels and association analysis for semen quality traits in Japanese men. *J Hum Genet* 2016;61:911–5.
- 27 World Health Organization. *WHO laboratory manual for the examination of human semen and sperm-cervical mucus interaction*. 4th edn. Cambridge: Cambridge University Press, 1999.
- 28 Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MA, Bender D, Maller J, Sklar P, de Bakker PI, Daly MJ, Sham PC. PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet* 2007;81:559–75.
- 29 Browning SR, Browning BL. Rapid and accurate haplotype phasing and missing-data inference for whole-genome association studies by use of localized haplotype clustering. *Am J Hum Genet* 2007;81:1084–97.
- 30 Browning BL, Browning SR. Genotype imputation with millions of reference samples. *Am J Hum Genet* 2016;98:116–26.
- 31 Auton A, Brooks LD, Durbin RM, Garrison EP, Kang HM, Korbel JO, Marchini JL, McCarthy S, McVean GA, Abecasis GR; 1000 Genomes Project Consortium. A global reference for human genetic variation. *Nature* 2015;526:68–74.
- 32 Sudmant PH, Rausch T, Gardner EJ, Handsaker RE, Abyzov A, Huddleston J, Zhang Y, Ye K, Jun G, Fritz MH, Konkel MK, Malhotra A, Stütz AM, Shi X, Casale F, Chen J, Hormozdiari F, Daya M, Chen K, Malig M, Chaisson MJP, Walter K, Meiers S, Kashin S, Garrison E, Auton A, Lam HYK, Mu XJ, Alkan C, Antaki D, Bae T, Cerveira E, Chines P, Chong Z, Clarke L, Dal E, Ding L, Emery S, Fan X, Gujral M, Kahveci F, Kidd JM, Kong Y, Lammeier EW, McCarthy S, Flicek P, Gibbs RA, Marth G, Mason CE, Menelaou A, Muzny DM, Nelson BJ, Noor A, Parrish NF, Pendleton M, Quidadamo A, Raeder B, Schadt EE, Romanovitch M, Schlattl A, Sebra R, Shabalina AA, Untergasser A, Walker JA, Wang M, Yu F, Zhang C, Zhang J, Zheng-Bradley X, Zhou W, Zichner T, Sebati J, Batzer MA, McCarroll SA, Mills RE, Gerstein MB, Bashir A, Stegle O, Devine SE, Lee C, Eichler EE, Korbel JO; 1000 Genomes Project Consortium. An integrated map of structural variation in 2,504 human genomes. *Nature* 2015;526:75–81.
- 33 Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ* 2003;327:557–60.
- 34 Pruim RJ, Welch RP, Sanna S, Teslovich TM, Chines PS, Glied TP, Boehnke M, Abecasis GR, Willer CJ. LocusZoom: regional visualization of genome-wide association scan results. *Bioinformatics* 2010;26:2336–7.
- 35 GTEx Consortium. The Genotype-Tissue Expression (GTEx) project. *Nat Genet* 2013;45:580–5.
- 36 Ward LD, Kellis M. HaploReg: a resource for exploring chromatin states, conservation, and regulatory motif alterations within sets of genetically linked variants. *Nucleic Acids Res* 2012;40:D930–4.
- 37 Boyle AP, Hong EL, Hariharan M, Cheng Y, Schaub MA, Kasowski M, Karczewski KJ, Park J, Hitz BC, Weng S, Cherry JM, Snyder M. Annotation of functional variation in personal genomes using RegulomeDB. *Genome Res* 2012;22:1790–7.
- 38 Zhang K, Cui S, Chang S, Zhang L, Wang J. i-GSEA4GWAS: a web server for identification of pathways/gene sets associated with traits by applying an improved gene set enrichment analysis to genome-wide association study. *Nucleic Acids Res* 2010;38:W90–5.
- 39 Marte BM, Jeschke M, Graus-Porta D, Taverna D, Hofer P, Groner B, Yarden Y, Hynes NE. Neu differentiation factor/hergulin modulates growth and differentiation of HC11 mammary epithelial cells. *Mol Endocrinol* 1995;9:14–23.
- 40 Yang Y, Spitzer E, Meyer D, Sachs M, Niemann C, Hartmann G, Weidner KM, Birchmeier C, Birchmeier W. Sequential requirement of hepatocyte growth factor and neuregulin in the morphogenesis and differentiation of the mammary gland. *J Cell Biol* 1995;131:215–26.
- 41 Carpenter G, Liao HJ. Trafficking of receptor tyrosine kinases to the nucleus. *Exp Cell Res* 2009;315:1556–66.
- 42 Roskoski R. The ErbB/HER family of protein-tyrosine kinases and cancer. *Pharmacol Res* 2014;79:34–74.
- 43 Sartor CI, Zhou H, Kozłowska E, Guttridge K, Kawata E, Caskey L, Harrelson J, Hynes N, Ethier S, Calvo B, Earp HS. Her4 mediates ligand-dependent antiproliferative and differentiation responses in human breast cancer cells. *Mol Cell Biol* 2001;21:4265–75.
- 44 Kim HC, Lee JY, Sung H, Choi JY, Park SK, Lee KM, Kim YJ, Go MJ, Li L, Cho YS, Park M, Kim DJ, Oh JH, Kim JW, Jeon JP, Jeon SY, Min H, Kim HM, Park J, Yoo KY, Noh DY, Ahn SH, Lee MH, Kim SW, Lee JW, Park BW, Park WY, Kim EH, Kim MK, Han W, Lee SA, Matsuo K, Shen CY, Wu PE, Hsiung CN, Lee JY, Kim HL, Han BG, Kang D. A genome-wide association study identifies a breast cancer risk variant in ERBB4 at 2q34: results from the Seoul Breast Cancer Study. *Breast Cancer Res* 2012;14:R56.
- 45 Day FR, Hinds DA, Tung JY, Stolk L, Styrkarsdottir U, Saxena R, Bjornes A, Broer L, Dunger DB, Halldorsson BV, Lawlor DA, Laval G, Mathieson I, McCardle WL, Louwers Y, Meun C, Ring S, Scott RA, Sulem P, Uitterlinden AG, Wareham NJ, Thorsteinsdottir U, Welt C, Stefansson K, Laven JS, Ong KK, Perry JR. Causal mechanisms and balancing

- selection inferred from genetic associations with polycystic ovary syndrome. *Nat Commun* 2015;6:8464.
- 46 Locke AE, Kahali B, Berndt SI, Justice AE, Pers TH, Day FR, Powell C, Vedantam S, Buchkovich ML, Yang J, Croteau-Chonka DC, Esko T, Fall T, Ferreira T, Gustafsson S, Kutalik Z, Luan J, Mägi R, Randall JC, Winkler TW, Wood AR, Workalemahu T, Faul JD, Smith JA, Zhao JH, Zhao W, Chen J, Fehrmann R, Hedman ÅK, Karjalainen J, Schmidt EM, Absher D, Amin N, Anderson D, Beekman M, Bolton JL, Bragg-Gresham JL, Buyske S, Demirkan A, Deng G, Ehret GB, Feenstra B, Feitosa MF, Fischer K, Goel A, Gong J, Jackson AU, Kanoni S, Kleber ME, Kristiansson K, Lim U, Lotay V, Mangino M, Leach IM, Medina-Gomez C, Medland SE, Nalls MA, Palmer CD, Pasko D, Pechlivanis S, Peters MJ, Prokopenko I, Shungin D, Stančáková A, Strawbridge RJ, Sung YJ, Tanaka T, Teumer A, Trompet S, van der Laan SW, van Setten J, Van Vliet-Ostaptchouk JV, Wang Z, Yengo L, Zhang W, Isaacs A, Albrecht E, Ärnlöv J, Arscott GM, Attwood AP, Bandinelli S, Barrett A, Bas IN, Bellis C, Bennett AJ, Berne C, Blagieva R, Blüher M, Böhringer S, Bonnycastle L, Böttcher Y, Boyd HA, Bruinenberg M, Caspersen IH, Chen YI, Clarke R, Daw EW, de Craen AJM, Delgado G, Dimitriou M, Doney ASF, Eklund N, Estrada K, Eury E, Folkersen L, Fraser RM, Garcia ME, Geller F, Giedraitis V, Gigante G, Go AS, Golay A, Goodall AH, Gordon SD, Gorski M, Grabe HJ, Grallert H, Grammer TB, Gräßler J, Grönberg H, Groves CJ, Haessler J, Hall P, Haller T, Hallmans G, Hartman CA, Hassinen M, Hayward C, Heard-Costa NL, Helmer Q, Hengstenberg C, Holmen O, Hottenga JJ, James AL, Jeff JM, Johansson Å, Jolley J, Juliusdottir T, Kinnunen L, Koenig W, Koskenvuo M, Kratzer W, Laitinen J, Lamina C, Leander K, Lee NR, Lichtner P, Lind L, Lindström J, Lo KS, Lobbens S, Lohrbein R, Lu Y, Mach F, Magnusson PKE, Mahajan A, McArdle WL, McLachlan S, Menni C, Merger S, Mihailov E, Milani L, Moayyeri A, Monda KL, Morken MA, Mulas A, Müller G, Müller-Nurasyid M, Musk AW, Nagaraja R, Nöthen MM, Nolte IM, Pilz S, Rayner NW, Renstrom F, Rettig R, Ried JS, Ripke S, Robertson NR, Rose LM, Sanna S, Schernagl H, Scholtens S, Schumacher FR, Scott WR, Seufferlein T, Shi J, Smith AV, Smolonska J, Stanton AV, Steinthorsdottir V, Stirrups K, Stringham HM, Sundström J, Swertz MA, Swift AJ, Syvänen AC, Tan ST, Tayo BO, Thorand B, Thorleifsson G, Tyrer JP, Uh HW, Vandenput L, Verhulst FC, Vermeulen SH, Verweij N, Vonk JM, Waite LL, Warren HR, Waterworth D, Weedon MN, Wilkens LR, Willenborg C, Wilsaard T, Wojczynski MK, Wong A, Wright AF, Zhang Q, Brennan EP, Choi M, Dastani Z, Drong AW, Eriksson P, Franco-Cereceda A, Gädin JR, Gharavi AG, Goddard ME, Handsaker RE, Huang J, Karpe F, Kathiresan S, Keildson S, Kiryluk K, Kubo M, Lee JY, Liang L, Lifton RP, Ma B, McCarrroll SA, McKnight AJ, Min JL, Moffatt MF, Montgomery GW, Murabito JM, Nicholson G, Nyholt DR, Okada Y, Perry JRB, Dorajoo R, Reinmaa E, Salem RM, Sandholm N, Scott RA, Stolk L, Takahashi A, Tanaka T, van 't Hooft FM, Vinkhuyzen AAE, Westra HJ, Zheng W, Zondervan KT, Heath AC, Arveiler D, Bakker SJL, Beilby J, Bergman RN, Blangero J, Bovet P, Campbell H, Caulfield MJ, Cesana G, Chakravarti A, Chasman DI, Chines PS, Collins FS, Crawford DC, Cupples LA, Cusi D, Danesh J, de Faire U, den Ruijter HM, Dominiczak AF, Erbel R, Erdmann J, Eriksson JG, Farrall M, Felix SB, Ferrannini E, Ferrières J, Ford I, Forouhi NG, Forrester T, Franco OH, Gansevoort RT, Gejman PV, Gieger C, Gottesman O, Gudnason V, Gyllenstein U, Hall AS, Harris TB, Hattersley AT, Hicks AA, Hindorf LA, Hingorani AD, Hofman A, Homuth G, Hovingh GK, Humphries SE, Hunt SC, Hyppönen E, Illig T, Jacobs KB, Jarvelin MR, Jöckel KH, Johansen B, Jousilahti P, Jukema JW, Jula AM, Kaprio J, Kastelein JJP, Keinanen-Kiukkaanniemi SM, Kiemeny LA, Knekt P, Koonen JS, Kooperberg C, Kovacs P, Kraja AT, Kumari M, Kuusisto J, Lakka TA, Langenberg C, Marchand LL, Lehtimäki T, Lyssenko V, Männistö S, Marette A, Matise TC, McKenzie CA, McKnight B, Moll FL, Morris AD, Morris AP, Murray JC, Nelis M, Ohlsson C, Oldehinkel AJ, Ong KK, Madden PAF, Pasterkamp G, Peden JF, Peters A, Postma DS, Pramstaller PP, Price JF, Qi L, Raitakari OT, Rankinen T, Rao DC, Rice TK, Ridker PM, Rioux JD, Ritchie MD, Rudan I, Salomaa V, Samani NJ, Saramies J, Sarzynski MA, Schunkert H, Schwarz PEH, Sever P, Shuldiner AR, Sinisalo J, Stolk RP, Strauch K, Tönjes A, Trégouët DA, Tremblay A, Tremoli E, Virtamo J, Vohl MC, Völker U, Waeber G, Willemsen G, Witteman JC, Zillikens MC, Adair LS, Amouyel P, Asselbergs FW, Assimes TL, Bochud M, Boehm BO, Boerwinkle E, Bornstein SR, Bottinger EP, Bouchard C, Cauchi S, Chambers JC, Chanock SJ, Cooper RS, de Bakker PIW, Dedoussis G, Ferrucci L, Franks PW, Froguel P, Groop LC, Haiman CA, Hamsten A, Hui J, Hunter DJ, Hveem K, Kaplan RC, Kivimäki M, Kuh D, Laakso M, Liu Y, Martin NG, März W, Melbye M, Metspalu A, Moebus S, Munroe PB, Njølstad I, Oostra BA, Palmer CNA, Pedersen NL, Perola M, Pérusse L, Peters U, Power C, Quertermous T, Rauramaa R, Rivadeneira F, Saaristo TE, Saleheen D, Sattar N, Schadt EE, Schlessinger D, Slagboom PE, Snieder H, Spector TD, Thorsteinsdottir U, Stumvoll M, Tuomilehto J, Uitterlinden AG, Uusitupa M, van der Harst P, Walker M, Wallaschofski H, Wareham NJ, Watkins H, Weir DR, Wichmann HE, Wilson JF, Zanen P, Borecki IB, Deloukas P, Fox CS, Heid IM, O'Connell JR, Strachan DP, Stefansson K, van Duijn CM, Abecasis GR, Franke L, Frayling TM, McCarthy MI, Visscher PM, Scherag A, Willer CJ, Boehnke M, Mohlke KL, Lindgren CM, Beckmann JS, Barroso I, North KE, Ingelsson E, Hirschhorn JN, Loos RJF, Speliotes EK, Åk H, As G, Å J, Ks L, Hw U, Study LC, Consortium A, Group A-BMIW; Lifelines Cohort Study ADIPOGen Consortium AGEN-BMI Working Group CARDIOGRAMplusC4D Consortium CKDGen Consortium GLGC ICBP MAGIC Investigators MuTHER Consortium MIGen Consortium PAGE Consortium ReproGen Consortium GENIE Consortium International Endogene Consortium. Genetic studies of body mass index yield new insights for obesity biology. *Nature* 2015;518:197–206.
- 47 Naillat F, Veikkolainen V, Miinalainen I, Sipilä P, Poutanen M, Elenius K, Vainio SJ. ErbB4, a receptor tyrosine kinase, coordinates organization of the seminiferous tubules in the developing testis. *Mol Endocrinol* 2014;28:1534–46.
- 48 Sandholm N, Salem RM, McKnight AJ, Brennan EP, Forsblom C, Isakova T, McKay GJ, Williams WW, Sadlier DM, Mäkinen VP, Swan EJ, Palmer C, Boright AP, Ahlqvist E, Deshmukh HA, Keller BJ, Huang H, Ahola AJ, Fagerholm E, Gordin D, Harjutsalo V, He B, Heikkilä O, Hietala K, Kytö J, Lahermo P, Lehto M, Lithovius R, Osterholm AM, Parkkonen M, Pitkäniemi J, Rosengård-Bärlund M, Saraheimo M, Sarti C, Söderlund J, Soro-Paavonen A, Syreeni A, Thorn LM, Tikkanen H, Tolonen N, Tryggvason K, Tuomilehto J, Wadén J, Gill GV, Prior S, Guiducci C, Mirel DB, Taylor A, Hosseini SM, Parving HH, Rossing P, Tarnow L, Ladenvall C, Alhenc-Gelas F, Lefebvre P, Rigalleau V, Rousset R, Tregouet DA, Mastroroni A, Mastroroni S, Falhammar H, Gu T, Möllsten A, Cimponeriu D, Ioana M, Mota M, Mota E, Serafinceanu C, Stavarachi M, Hanson RL, Nelson RG, Kretzler M, Colhoun HM, Panduru NM, Gu HF, Brismar K, Zerbini G, Hadjadj S, Marre M, Groop L, Lajer M, Bull SB, Waggott D, Paterson AD, Savage DA, Bain SC, Martin F, Hirschhorn JN, Godson C, Florez JC, Groop PH, Maxwell AP; DCCT/EDIC Research Group. New susceptibility loci associated with kidney disease in type 1 diabetes. *PLoS Genet* 2012;8:e1002921.
- 49 Liu SY, Zhang CJ, Peng HY, Sun H, Lin KQ, Huang XQ, Huang K, Chu JY, Yang ZQ. Strong association of SLC1A1 and DPFF3 gene variants with idiopathic male infertility in Han Chinese. *Asian J Androl* 2017;19:486–92.