ORIGINAL ARTICLE

Genetic linkage analysis of a large family identifies *FIGN* as a candidate modulator of reduced penetrance in heritable pulmonary arterial hypertension

Pau Puigdevall,¹ Lucilla Piccari,² Isabel Blanco,^{2,3} Joan Albert Barberà,^{2,3} Dan Geiger,⁴ Celia Badenas,^{2,5} Montserrat Milà,^{2,5} Robert Castelo,^{1,6} Irene Madrigal^{2,5}

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/jmedgenet-2018-105669>).

¹Department of Experimental and Health Sciences, Universitat Pompeu Fabra, Barcelona, Spain

²Hospital Clínic de Barcelona, Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS), Barcelona, Spain

³Centro de Investigación Biomédica en Red de Enfermedades Respiratorias (CIBERES), Madrid, Spain

⁴Faculty of Computer Science, Technion Israel Institute of Technology, Haifa, Israel

⁵Centro de Investigación Biomédica en Red de Enfermedades Raras (CIBERER), Instituto de Salud Carlos III, Madrid, Spain

⁶Institut Hospital del Mar d'Investigacions Mèdiques (IMIM), Barcelona, Spain

Correspondence to

Dr Robert Castelo, Department of Experimental and Health Sciences, Universitat Pompeu Fabra, Barcelona 08002, Spain; robert.castelo@upf.edu

Received 8 August 2018

Revised 12 February 2019

Accepted 16 February 2019

Published Online First 20 March 2019

© Author(s) (or their employer(s)) 2019. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Puigdevall P, Piccari L, Blanco I, et al. *J Med Genet* 2019;**56**:481–490.

ABSTRACT

Background Mapping the genetic component of molecular mechanisms responsible for the reduced penetrance (RP) of rare disorders constitutes one of the most challenging problems in human genetics. Heritable pulmonary arterial hypertension (PAH) is one such disorder characterised by rare mutations mostly occurring in the bone morphogenetic protein receptor type 2 (*BMPR2*) gene and a wide heterogeneity of penetrance modifier mechanisms. Here, we analyse 32 genotyped individuals from a large Iberian family of 65 members, including 22 carriers of the pathogenic *BMPR2* mutation c.1472G>A (p.Arg491Gln), 8 of them diagnosed with PAH by right-heart catheterisation, leading to an RP rate of 36.4%.

Methods We performed a linkage analysis on the genotyping data to search for genetic modifiers of penetrance. Using functional genomics data, we characterised the candidate region identified by linkage analysis. We also predicted the haplotype segregation within the family.

Results We identified a candidate chromosome region in 2q24.3, 38 Mb upstream from *BMPR2*, with significant linkage (LOD=4.09) under a PAH susceptibility model. This region contains common variants associated with vascular aetiology and shows functional evidence that the putative genetic modifier is located in the upstream distal promoter of the fidgetin (*FIGN*) gene.

Conclusion Our results suggest that the genetic modifier acts through *FIGN* transcriptional regulation, whose expression variability would contribute to modulating heritable PAH. This finding may help to advance our understanding of RP in PAH across families sharing the p.Arg491Gln pathogenic mutation in *BMPR2*.

INTRODUCTION

Pulmonary arterial hypertension (PAH) is a rare disease characterised by an abnormal rise in mean pulmonary arterial pressure (≥ 25 mm Hg at rest), which leads to a progressive increase in pulmonary vascular resistance and ultimately to death, due to right ventricular failure.¹ From 2% to 4% of PAH patients suffer from heritable pulmonary arterial hypertension (HPAH), with an overall HPAH prevalence below one case per million adults.^{2,3} PAH was initially described by Dresdale *et al*,⁴ and its hereditary subtype is defined by either the presence

of a known genetic defect linked to the disease or by a positive family history. Despite substantial progress in diagnosing HPAH, long-term survival for most patients is extremely poor.

Mutations in the bone morphogenetic protein receptor type 2 gene (*BMPR2*),^{5,6} a transforming growth factor beta (TGF- β) superfamily member, have been detected in 75%–80% of PAH cases.⁷ Other genes have been implicated in the disease, mainly involving the TGF- β signalling pathway (*ALK1*,⁸ *ENG*⁹ and *SMAD8*¹⁰), as well as other genes (*CAV1*,¹¹ *KCNK3*¹²). The molecular mechanisms by which *BMPR2* mutations cause the disease are still not fully understood. HPAH is inherited as an autosomal dominant disease. However, not all *BMPR2* mutation carriers develop the disease, highlighting the presence of reduced penetrance (RP). Consequently, a primary *BMPR2* mutation is necessary, but not sufficient by itself, to cause PAH. The penetrance of PAH among *BMPR2* mutation carriers is estimated to be around 20%.¹³

There are several proposed mechanisms to explain RP¹⁴ in human disease, including genetic modifiers, the molecular context of mutations, patient characteristics such as age or sex, or environmental conditions. In the case of PAH, *BMPR2* mutations alter multiple pathways supporting the hypothesis that additional genetic factors may confer susceptibility onto *BMPR2* carriers.¹⁵ In recent years, many efforts have thus focused on finding those genetic modifiers.^{16–21} However, most of these studies identify different genetic mechanisms, supporting a wide heterogeneity among the acting modifier loci. This suggests that new modifiers are likely to be found as more affected families are investigated.

We studied a large Iberian family (n=65 subjects, five generations) affected by PAH and segregating with the *BMPR2* missense mutation p.Arg491Gln (rs137852749, c.1472G>A).²² The penetrance of PAH among *BMPR2* mutation carriers is 36.4%. Sequence-level characteristics may contribute to this increased penetrance because this particular variant occurs in a highly conserved kinase domain position of a superfamily type II TGF- β receptor.²³ This mutation leads to a dominant-negative inhibition effect on the wild-type receptor function, altering the *BMPR2* heteromerisation

and impacting the selection of specific signalling pathways downstream.²⁴

METHODS

Subjects and phenotypic characterisation

The family studied comprises 65 Spanish individuals spanning five different generations.²² Eight of the individuals were diagnosed with PAH, confirmed by right-heart catheterisation, showing a mean pulmonary arterial pressure ≥ 25 mm Hg. DNA extractions and mutational studies of the *BMPR2* gene from whole blood were carried out following standard procedures.²⁵ The pathogenic *BMPR2* mutation c.1472G>A (p.Arg491Gln)²³ segregated with the disease in an autosomal dominant mode of inheritance (MOI). All 8 PAH subjects carried the *BMPR2* mutation, while the other 14 carriers were healthy at the time of examination. The remaining individuals were reported as healthy non-carriers, except for eight individuals whose disease and/or genetic status were unknown, including four obligate carriers due to carrier offspring and suspicion of PAH, for which haemodynamic confirmation was not available (S01, S06, S08 and S10). Living subjects carrying the mutation within the family are regularly followed up by their specialists of reference.

Genotyping and data preprocessing

We genotyped 32 individuals selected from the family being studied, with the Illumina microarray chip *Infinium Core-Exome-24 BeadChip v1.1* ($n=5\,51839$ SNPs). Genotype calling and sample clustering were performed with the *Illumina Bead-Studio* software. Five individuals presented a genotyping call rate <0.99 (labelled in grey: T25, T26, T27, T28 and T29) and, along with four non-genotyped relatives (S12, S13, S30 and S33), were discarded from further analysis, leaving 56 out of the initial 65 members.

We filtered out autosomal and X-linked variants that were not biallelic SNPs, as well as biallelic SNPs with either missing genotypes in more than six samples or that could not be unambiguously annotated to dbSNP Build 144. We also removed variants that lacked the genetic and physical position provided by Illumina, as well as population allele frequency (AF) data from the ExAC²⁶ release 0.3 or the 1000 Genomes Project,²⁷ Phase 3. All reported genomic coordinates used the GRCh37/hg19 genome version as reference.

The PLINK²⁸ Mendelian error tool was also applied to the dataset to remove segregation inconsistencies. For chromosome X, an additional filter discarded all variants containing heterozygous genotypes in men. For pseudoautosomal regions (XY), only variants falling within the PAR1 region were considered. Within that region, the Rutgers (v3) genetic sex-specific map²⁹ was used instead of the Illumina coordinates. Following these preprocessing steps, 491 896 SNPs remained in the dataset (480 028 autosomal, 11 759 from chromosome X and 73 from XY-PAR1 region; see online figure S1).

Linkage analysis

We performed genome-wide linkage analysis with the previously filtered SNPs. To assess the robustness of the observed results, linkage analysis was conducted using five independent software programmes based on two-point (Mendel,³⁰ Pseudomarker³¹ and Superlink³²) and multipoint (Merlin³³ and Morgan³⁴) linkage approaches (online figure S1).

The software Mendel³⁰ used all the filtered variants as input, while Pseudomarker³¹ was unable to process variants in the PAR1 region. Mendel ran a two-point linkage under the

Location_scores option using two flanking points. Pseudomarker was run using the runPM programme (see Web resources), which internally removes all the monomorphic variants present in the input. Superlink³² reduced the number of input variants based on various clustering, cleaning and filtering algorithms.

For multipoint linkage, we first ran parametric and non-parametric tests in Merlin.³³ We were forced to prune the family pedigree (67 bits) due to the complexity upper bound of the Lander-Green algorithm implementation in Merlin (24 bits). Accordingly, we selected a subtree of the family pedigree that maximised the statistical power of linkage, as follows. We first prioritised the selection of genotyped affected carriers, and then included as many genotyped healthy carriers as possible. Finally, we included genotyped healthy non-carrier individuals (online figure S2). The input number of SNPs was drastically reduced by a linkage disequilibrium (LD) filtering performed with PLINK 1.07 ($n=33\,378$ SNPs), using a window size of 10 SNPs, a sliding window of 5 SNPs and a variance inflation factor threshold of 2 units. We also incorporated the ExAC minor allele frequencies (MAF) into the model. Where these values were not available, we used MAF data from the 1000 Genomes Project. We also ran multipoint linkage with the whole pedigree using Superlink,³² restricted to SNPs within chr2:161–167 Mb. Given the size of the pedigree, we could only run non-overlapping windows of a maximum size of five SNPs. Because calculations are sensitive to the subset of SNPs within the sliding window, we performed five different runs beginning at each of the five possible starting SNPs of the analysed region.

The software Morgan³⁴ was also used to estimate multipoint LOD scores. This programme is not restricted by pedigree complexity, as it uses the Markov Chain Monte Carlo algorithm, although the number of SNPs is computationally limiting. Consequently, we ran an analysis using a window of 300 SNPs on the regions of interest highlighted by Merlin.

We began the linkage analysis by verifying, as a quality control, the presence of significant linkage at the *BMPR2* locus selecting *BMPR2* mutation carriers as affected individuals. We then searched for a linkage signal from another locus with an independent contribution to the disease, whose genotype differentiates affected carriers from the rest of the individuals (online Supplemental methods). Finally, we considered a susceptibility model that could explain the low penetrance in HPAH (online figure S3). Under this model, all affected *BMPR2* carriers possess a genetic modifier that increases the risk of the disease as part of a two-hit mechanism, while healthy non-carriers could have the modifier without altering their risk of PAH.

Under a susceptibility model, there are four different scenarios, which depend on the AF (common or rare in the population) and the MOI (dominant or recessive). We discarded two of these scenarios because they were extremely unlikely (rare/recessive, common/dominant). Another scenario (rare/dominant) was discarded because it assumed a two-hit occurrence of rare variants. Consequently, the parametric model for the susceptibility tests was built using a common AF ($d=0.999$) and a recessive MOI (see online Supplemental methods). We also permitted the existence of locus heterogeneity by defining a low phenocopy rate of 2%.

The age at disease onset is a particular confounder for the clinical condition as healthy carriers can develop the condition later in life and thus switch to affected carriers. The standard approach to control for this age-dependent penetrance effect in linkage analysis is through liability classes. However, the age-of-onset exhibits a very large variability, ranging from 5 to 75 years, and appears to be affected by genetic anticipation. This makes

Figure 1 Family tree with HPAH disease cosegregation. Identifiers of genotyped individuals begin with the capital letter T, while non-genotyped begin with S. Solid black dots indicate affected mutation carriers while vertically striped dots indicate asymptomatic carriers and open dots correspond to healthy non-carriers. Grey dots indicate individuals of unknown phenotype and/or genotype while crossed dots correspond to deceased individuals. S01, S06, S08 and S10 individuals are obligate carriers due to carrier offspring. HPAH, heritable pulmonary arterial hypertension .

a general penetrance function more appropriate than liability classes.

The individuals most at-risk of developing the condition are the youngest healthy carriers. Within our family, three healthy carriers are <10 years of age (T20, T21 and T24; online table S1). Given this uncertainty, we ran several linkage tests including all the family (or the fraction allowed by Merlin) and the seven pedigree combinations created by leaving aside the three youngest healthy carriers. This included the removal of only one individual at a time (three combinations), each pair of individuals (three combinations) and excluding all three at once (one combination). We chose this strategy to avoid direct speculation on individual phenotypes and due to the limited number of young individuals to build liability classes based on the age of onset.

RESULTS
Clinical evaluation of *p.Arg491Gln* mutation carrier individuals

We studied a large multiplex family composed of 65 members (figure 1), including 22 carriers of the pathogenic *BMPR2* mutation c.1472G>A (*p.Arg491Gln*). At the time of the examination, eight of these individuals were affected. Table 1 shows the clinical and haemodynamic characteristics of healthy and affected carriers. Unfortunately, the available data do not allow us to compare the haemodynamic trajectories between healthy and affected carriers for several reasons. First, affected carriers have no pulmonary arterial pressure (systolic) (PAPs) echocardiogram values because they were diagnosed as PAH symptoms occurred, with the exception of two individuals. Second, the currently healthy carriers who were screened by echocardiogram either did not show the tricuspid regurgitation needed to measure PAPs values or data were not available. While pulmonary arterial pressure (mean) (PAPm) values are available for affected carriers, only one healthy carrier with suspected PAH, underwent right-heart catheterisation (T29). As expected, the haemodynamic values (PAPs, PAPm and pulmonary vascular resistance) of this healthy carrier were very different from the average values among affected carriers.

Table 1 also shows that females are more affected than males, as previously reported.²² This difference is also observed when comparing the penetrance of female (45.5%) to male carriers (27.3%). The age of onset in this family is highly variable, spanning from 5 to 75 years (figure 2), decreasing with each new

generation. More concretely, in generation II, PAH was diagnosed in one patient when he was 75.4 years old, in generation III at 47.1±5.2 years (n=3) and in generation IV at 18.16±12.58 years (n=4); see figure 2. This pattern is consistent with genetic anticipation, although clinical improvements in the diagnosis and earlier tracking of younger carriers may confound this observation.

Another mechanism that could explain the RP is genomic imprinting (online Figure S4; see Supplemental Methods). Despite the differences observed in the penetrance of carrier individuals with paternal transmission (f; 50%) and maternal transmission (m; 25%), the parental origin of the mutation does not entirely explain the phenotype of the progeny.

Table 1 Clinical and haemodynamic characteristics of carriers of the *BMPR2* *p.Arg491Gln* mutation

	Healthy carriers (n=14)	Affected carriers (n=8)
Sex (female): n (%)	6 (42.9)	5 (62.5)
Years free of PAH: mean±SD	45.3±26.8	36.2±23.1
PAPs echocardiogram (mm Hg): mean±SD	NA*	90±0 (n=2)†
PAPs right heart catheterisation (mm Hg): mean±SD	23 (n=1)‡	81±14.8 (n=7)§
PAPm (mm Hg): mean±SD	17 (n=1)‡	52±10 (n=7)¶
CI (L/min/m ²): mean±SD	2.23 (n=1)‡	2.05±0.88 (n=6)¶
PAWP (mm Hg): mean±SD	12 (n=1)‡	9±3 (n=5)¶
PVR (WU): mean±SD	1.54 (n=1)‡	15±7.94 (n=4)¶

*No PAPs values could be measured by echocardiogram for five screened healthy carriers due to lack of tricuspid regurgitation. The remaining nine healthy carriers were not screened, or had no data available.

†The only two affected carriers with echocardiogram PAPs values were measured immediately prior to the diagnosis by right-heart catheterisation.

‡Individual T29 was the only healthy carrier to undergo right-heart catheterisation.

§All the eight affected individuals were diagnosed with a PAPm ≥25 mm Hg after right-heart catheterisation. This abnormal measure was confirmed for all the individuals by their specialist of reference, although we lacked the PAPm-specific value for one of the affected patients (S18).

¶The haemodynamic parameters (CI, PAWP, PVR) were determined by right-heart catheterisation for all the affected individuals, although for some of these, the actual values are not available.

PAPs, pulmonary arterial pressure (systolic); PAPm, pulmonary arterial pressure (mean); CI, cardiac index; PAWP, pulmonary arterial wedge pressure ; PVR, pulmonary vascular resistance.

J Med Genet: first published as 10.1136/jmedgenet-2018-105669 on 20 March 2019. Downloaded from <http://jmg.bmj.com/> on April 26, 2023 by guest. Protected by copyright.

PAH susceptibility among *BMPR2* p.Arg491Gln mutation carriers is significantly linked to region 2q24.2-q24.3

We then considered the age-dependent penetrance effect on the youngest healthy carriers (T20, T21 and T24) by trimming one, two or three of those subjects at a time from the pedigree. The results for these additional tests revealed a variant at 2q24.2 (rs17716942) with significant two-point linkage for the same two combinations of individuals using both Mendel (online figure S12C and G) and Pseudomarker (online figure S13C and G). Those two combinations correspond to the trimming of T21 (max LOD=4.14), and of the pair T21–T24 (max LOD=3.86). Additionally, Pseudomarker reported other variants with weaker (but still significant) linkage, although Mendel did not reproduce them. This is also the case for variant rs6436140 detected at 2q35 (online figure S13A, C, D and G).

statistics coupled with the Kong and Cox exponential method) did not support the candidate region. However, the regions detected using the parametric method with two-point and multipoint linkage clearly overlap at the same region, 2q24.2 (figure 3A).

These results suggest that the region 2q24.2-q24.3 is a strong candidate to harbour a genetic modifier that modulates HPAH susceptibility. We identified 12 annotated protein-coding genes within this region (figure 3B). Among them, *FIGN* was found to be one of the best candidates, as the local maximum score of the candidate region overlaps a variant (rs12692701) from the intron 2 of the annotated isoform NM_018086.3 of this gene. Interestingly, in the region of maximum linkage (LOD=4.09), several SNPs are associated with blood pressure according to the traits reported in the genome-wide association study (GWAS) Catalog.

The candidate susceptibility region (2q24.2-q24.3) is enriched by SNPs from the GWAS Catalog associated with blood pressure

We analysed the functional impact of the 529 SNPs located within the candidate region of LOD >3.3 and in LD with annotated SNPs from the GWAS Catalog.³⁵ More concretely, we performed an enrichment analysis of these SNPs with respect to all the SNPs in the human genome for which we could map experimental factor ontology (EFO) annotations from the GWAS Catalog; see online Supplemental Methods. We observed 24 EFO terms that were significantly enriched by SNPs in the candidate region (adjusted Holm $p < 10^{-3}$, Size >5 , Count >5 , OR >2 , one-tailed Fisher's exact test) when compared with the entire GWAS Catalog (online table S2). Interestingly, EFO terms associated with cardiorespiratory phenotypes were significantly enriched (EFO:0004325: blood pressure, EFO:0006335: systolic blood pressure, EFO:0005763: pulse pressure measurement, EFO:0006995: response to diisocyanate and EFO:0000270: asthma), reinforcing the hypothesis of a potential contribution of that region to blood pressure.

When conducting the same analysis on the subregion with strongest linkage (LOD=4.09; [figure 3B](#)), 15 EFO terms showed as significant ([table 2](#)). Among them, the aforementioned cardiorespiratory phenotypes remained on the list. Seven GWAS SNPs, located upstream from the *FIGN* gene, were responsible for this enrichment: rs16849225,^{36 37} rs11891401³⁸ and rs1446468³⁹ for systolic blood pressure; rs13002573⁴⁰ and rs16849211³⁹ for pulse pressure measurement; rs76833157 and rs7576072 for both asthma and response to diisocyanate.⁴¹ Furthermore, one variant from a dbGAP study was associated with tunica media wall thickness of carotid arteries (rs1370493, pha003034-phs000221).⁴² These enriched EFO terms refer to systemic cardiopulmonary vasculature, and not to the specific pulmonary vasculature, which would be the most relevant for PAH physiology but which is unfortunately absent from the GWAS Catalog and EFO term annotations.

Conversely, the enrichment of cardiorespiratory phenotypes was not observed when selecting SNPs within the boundaries of the *FIGN* gene (online table S3). Upstream of *FIGN*, partially within the region of strongest linkage, a long intergenic non-coding RNA (lincRNA), *ENSG00000237844*, was exclusively enriched by cardiorespiratory phenotypes (online table S4). However, this lincRNA shows no expression in

Figure 3 Linkage analysis results. (A) LOD score profiles from different linkage analysis software in the genomic region between the highest LOD and the *BMPR2* locus. The profile Merlin-subpedALL was calculated from the subpedigree shown in online figure S1. Merlin-npl-exp is the non-parametric linkage with the same individuals as in Merlin-subpedALL. Profiles marked with 'no-Tx' were calculated excluding the 'Tx' individuals. In the case of Mendel and Pseudomarker, only SNPs with significant LOD score are shown. Except for Morgan, all programs highlight a region of significant linkage with the combinations that remove the healthy carriers T21 and T24. (B) LOD score profile from Merlin-noT21 in the 7 Mb region of significant LOD. The SNPs track shows the SNPs used by Merlin in the parametric multipoint linkage analysis. The genome-wide association study (GWAS) track shows SNPs from the GWAS Catalog that are associated with blood pressure (rs13002573, rs1446468, rs16849225 and rs11891401). The shaded region highlights SNPs with the strongest linkage (LOD=4.09), which includes the *FIGN* gene.

almost all tissues from the GTEx project,⁴³ excluding a mechanism mediated by this RNA molecule.

Common variation in the region of maximum linkage (2q24.3) impacts regulatory elements that potentially act as a distal promoter of the *FIGN* gene

We examined whether the seven GWAS SNPs associated with cardiorespiratory traits in the region of maximum linkage

overlap with regulatory elements upstream from the *FIGN* gene. We refer to this set of SNPs as candidate regulatory SNPs (CRS).

Hi-C data⁴⁴ from the IMR-90 cell line derived from human fetal lung tissue show that both these CRSs and the *FIGN* gene are located within the same topologically associated domain (figure 4A). In GTEx data, three of the CRSs (rs13002573, rs16849225 and rs16849211) are cis-eQTLs of the *FIGN* gene in aortal artery, oesophagus mucosa, skeletal muscle and

Table 2 Functional enrichment analysis on the region of maximum linkage. EFO terms enriched by 137 SNPs from the region with LOD=4.09 (chr2:163738883–165107298). EFO terms associated with cardiorespiratory traits are highlighted in bold

	EFO term identifier	Raw p value	OR	Expected count	Count	Size	EFO term description	Parental term cluster
1	EFO:0007677	1.07e–08	78.07	0.07	5	123	LDL peak particle diameter measurement	LDL peak particle diameter measurement
2	EFO:0007678	1.07e–08	78.07	0.07	5	123	Total fat intake measurement	Total fat intake measurement
3	EFO:0006335	4.10e–79	71.21	1.52	60	2694	Systolic blood pressure	Vital signs
4	EFO:0004325	1.41e–65	41.42	2.56	60	4553	Blood pressure	Vital signs
5	EFO:0003884	2.16e–20	35.73	0.55	17	978	Chronic kidney disease	Kidney disease
6	EFO:0005763	5.48e–24	33.01	0.76	21	1348	Pulse pressure measurement	Pulse pressure measurement
7	EFO:0004303	6.79e–60	32.90	3.20	60	5689	Vital signs	Vital signs
8	EFO:0006995	3.43e–16	30.42	0.52	14	921	Response to diisocyanate	Process
9	GO:0097332	3.11e–23	30.28	0.83	21	1467	Response to antipsychotic drug	Process
10	EFO:0003086	3.95e–12	10.82	1.78	17	3162	Kidney disease	Kidney disease
11	EFO:0000270	3.84e–08	7.22	2.13	14	3793	Asthma	Bronchial disease
12	EFO:1002018	5.90e–08	6.96	2.21	14	3929	Bronchial disease	Bronchial disease
13	GO:0050896	1.56e–09	3.72	12.01	36	21 335	Response to stimulus	Process
14	GO:0008150	6.59e–09	3.09	19.28	46	34 266	Biological process	Process
15	BFO:0000007	9.78e–09	3.04	19.52	46	34 697	Process	Process

pancreas tissues (figure 4B). Although none of these GTEx tissues correspond to pulmonary microvascular endothelial cells (PMVCs)—the tissue most relevant to PAH—these cis-eQTLs suggest the presence of molecular regulatory mechanisms that may be shared across tissues. The three CRSs are also located in the same LD block according to the web tool HaploReg,⁴⁵ which shows that some of the linked SNPs alter regulatory motifs and overlap enhancer histone marks (online figure S17). Chromatin state segmentation from the Roadmap Epigenomics Mapping Consortium (REMC) in the IMR-90 cell line and lung tissue shows changes between quiescent, heterochromatic, zinc finger and flanking promoter states across the LD block (online figure S18).

One of the linked SNPs in the LD block (rs4667728) is a cis-eQTL of *FIGN* (figure 4C). According to CENTIPEDE annotations⁴⁶ (online figure S18), this common variant affects protein binding in an open chromatin region with evidence of REMC data from DNaseI of a fetal lung donor. Moreover, this same variant overlaps with a DNA methylation signal from REMC bisulfite sequencing data of the IMR-90 cell line (figure 4D). The altered motif, predicted by CENTIPEDE, derives from the DREB1B factor, a transcriptional activator in *Arabidopsis thaliana* whose DNA-binding domain superfamily (IPR016177) is, according to the InterPro database, homologous to the methyl-CpG DNA binding (MBD) domain (IPR001739). MBD-containing proteins, such as MeCP2, MBD1 and MBD2 preferentially bind to methylated CpG, such as that overlapped by the rs4667728 SNP, whose reference allele is the corresponding cytosine.

To investigate whether the transcriptional regulation of *FIGN* is associated with PAH expressivity, we have analysed publicly available gene expression data from a study⁴⁷ that collected samples derived from the lung tissue of PAH patients and from the normal tissue of cancer patients undergoing surgical lobectomy. In a differential expression analysis between these two groups of samples, we found that *FIGN* is significantly overexpressed (1.3-fold) in PAH patients, after correcting for multiple testing at 1% FDR (online figure S19). We also found that *FIGN* is inversely and significantly co-expressed with *BMPR2* ($p=0.02$), using a linear model that adjusts for the PAH effect

in *FIGN* expression (online figure S19); see online Supplemental Methods.

All linked SNPs in the LD block, including rs4667728, have a common MAF of around 0.22 in the European population of the 1000 Genomes Project (online figure S17). The linkage signal and the described functional genomics evidence thus both support the hypothesis that common variation in the candidate susceptibility region (2q24.3) may impact regulatory elements acting as a distal promoter (250–500 kb) of the *FIGN* gene, whose expression would contribute to modulating HPAH penetrance.

Haplotype prediction

Predicted haplotypes in chromosome 2 were consistent with the *BMPR2* carrier status (online figure S20; see online Supplemental Methods). We observed that no healthy carriers were predicted to have a recombination between the *FIGN* (first candidate region) and *BMPR2* loci, except for T21. However, half of the affected carriers in the subpedigree (T14 and T11) do present such a recombination. Consistent with these observations, the variant with highest LOD score in two-point linkage (rs17716942) was heterozygous among all healthy carriers, except for T21, and homozygous in half of the affected carriers in the sub-pedigree (T14 and T11).

DISCUSSION

The region defined by the SNPs with LOD >3.3 in multipoint linkage is annotated with 12 protein-coding genes; among these, the *FIGN* gene was the only gene overlapping the markers with the highest LOD score. Moreover, one intronic variant (rs12692701) supports a direct linkage to the gene. However, while this region is functionally enriched with SNPs associated with cardiorespiratory phenotypes, these are not located within the gene boundaries of *FIGN*, but in its promoter region. Concretely, multiple sources of functional genomics data—GTEx cis-eQTLs, Hi-C interactions and REMC DNaseI and methylation data—support the hypothesis that some of these

Figure 4 Common variation upstream from *FIGN* contributes to its regulation and its role as HPAH penetrance modulator. (A) Hi-C data from the human IMR-90 cell line derived from fetal lung tissue. Vertical yellow stripes highlight SNPs from the GWAS catalogue associated with cardiorespiratory traits, which fall within the same topologically associated domain of the *FIGN* gene. (B) GTEx *FIGN* cis-eQTLs located at the region of maximum linkage (LOD=4.09). The first four tracks show raw p values, in logarithmic scale, from eQTLs overlapping SNPs associated with cardiorespiratory traits in the GWAS Catalog. The shaded region highlights an enrichment of eQTLs within a linkage disequilibrium (LD) block that includes three candidate regulatory SNPs (CRS: rs16849225, rs16849211, rs13002573) associated with systolic blood pressure and pulse pressure measurement. (C) Genetic effect on gene expression of *FIGN* for the cis-eQTL in the rs4667728 SNP. (D) Functional genomics evidence within the LD block of *FIGN* cis-eQTLs. The vertical yellow stripe highlights SNP rs4667728 predicted by CENTIPEDE annotations to alter protein binding by integrating REMC DNaseI data with protein DNA-binding motifs. This variant also overlaps a REMC methylation signal from the same IMR-90 cell-line as the DNaseI data. HPAH, heritable pulmonary arterial hypertension.

SNPs may contribute to the regulation of *FIGN* expression in relevant tissues for vascular aetiology.

Mukherjee *et al.*⁴⁸ characterised *FIGN* as a microtubule severing enzyme as well as a depolymerase, predominantly involved in mitosis but also contributing to cell migration and neuronal development. *FIGN* controls the conserved localisation of microtubules to centrosomes, with a crucial role in avoiding the formation of aneuploid or polyploid cells. The same study observed that the depletion of *FIGN* resulted in alterations in the structure of mitotic centrosomes. However, Wang *et al.*⁴⁹ have recently identified a *FIGN* intronic variant (rs2119289) that has a protective role against congenital heart disease. This finding is

compatible with the role of *FIGN* as contributing to modulating HPAH penetrance.

The mere existence of phenocopies in the susceptibility model implicitly assumes that HPAH may also be caused by genetic modifiers other than that which we found in the candidate region. Such locus heterogeneity in HPAH has already been reported by many different studies. These include the alteration of *BMPT2*^{15 16} and *CBLN2*¹⁷ gene expression, second-hit mechanisms of *BMPT2* with its promoter¹⁹ or with *EIF2AK4* mutations,¹⁸ as well as somatic chromosome abnormalities in the affected lungs.²⁰ These multiple findings support an oligogenic architecture of HPAH.

In summary, we found the *FIGN* gene to be a candidate modulator for RP in HPAH within the largest family reported to date segregating with the *BMPR2* pathogenic mutation c.1472G>A (p.Arg491Gln). Multiple sources of functional genomics data strongly support the most likely underlying genetic mechanism to be located in a regulatory region upstream from the *FIGN* gene, forming part of its distal promoter and modulating *FIGN* expression through common variation. A similar mechanism has been observed for familial neuroblastoma, in which genetic modifiers of penetrance for a pathogenic mutation in the *ALK* gene are identified within the same chromosome arm 2q.⁵⁰

Our study has several limitations and future research will be required to replicate and confirm the findings reported. First, we were unable to conduct a genome-wide analysis of the entire pedigree at once with currently available multipoint linkage analysis tools due to the algorithmic constraints on large families. Second, microarray genotyping data are sparse and whole-genome sequencing (WGS) data will be required to identify the specific variant, or variants, that act as genetic modifiers of HPAH within the candidate region. WGS data will also be useful to better resolve the individual haplotypes from chromosome 2 in mutation carrier individuals, where long-read sequencing technology would be the most reliable option. Third, we could not obtain PMVCs, the PAH target tissue, from carrier individuals in the family. This tissue would allow us to profile the most relevant RNA and protein expression, which could help to characterise the molecular mechanism triggered by the proposed genetic modifier. Finally, further functional in vitro and in vivo assays will be required to validate this molecular mechanism.

Web resources

runPM: <http://www.jurgott.org/linkage/runPM.html>

GTE Portal: <https://www.gtexportal.org>

Correction notice This article has been corrected since it was published online first. A technical error in the PDF (figure 3 was not displaying) has been corrected.

Acknowledgements This study was supported by the Spanish Instituto de Salud Carlos III (ISCIII) [PI15/00483], co-financed by Fondo Europeo de Desarrollo Regional (FEDER) 'una manera de hacer Europa', Catalan AGAUR [SGR17-1134; SGR17-1020; FI-DGR 2015], Spanish MINECO/FEDER (TIN2015-71079-P) and the 'Maria de Maeztu Unit of Excellence' (MDM-2014-0370). The 'CIBER de Enfermedades Raras' is an initiative of the ISCIII. We want to thank the 'CERCA Programme' from the Autonomous Catalan Government. We also thank Manuel López-Meseguer and Luis Borderías for providing clinical data, and Roger Piqué-Regi for assistance in using CENTIPEDE. We acknowledge the NCBI database of Genotypes and Phenotypes (dbGaP, <http://www.ncbi.nlm.nih.gov/gap>). Controlled-access genotype and phenotype data were obtained following authorisation by the appropriate Data Access Committee to GTEx data (dbGaP Study Accession number phs000424.v7.p2).

Contributors IM, MM, CB and RC designed the project. IM, MM, CB, LP and JAB collected the data. PP, DG and RC analysed the data. PP, CB, IM, LP and RC drafted the manuscript. All authors revised and approved the manuscript.

Funding This work was supported by the Spanish Instituto de Salud Carlos III (ISCIII) [PI15/00483], co-financed by Fondo Europeo de Desarrollo Regional (FEDER) 'una manera de hacer Europa', Catalan AGAUR [SGR17-1134; SGR17-1020; FI-DGR 2015], Spanish MINECO/FEDER (TIN2015-71079-P) and the 'Maria de Maeztu Unit of Excellence' (MDM-2014-0370). The 'CIBER de Enfermedades Raras' is an initiative of the ISCIII. We want to thank the "CERCA Programme" from the Autonomous Catalan Government.

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval The study was approved by the ethical committee of the Hospital Clínic de Barcelona. All participants provided written, informed consent.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement Genotype and phenotype data reported in this paper are available through the European Genome-phenome Archive (EGA) under accession number EGAS00001003123.

REFERENCES

- Galiè N, Humbert M, Vachieri J-L, Gibbs S, Lang I, Torbicki A, Simonneau G, Peacock A, Vonk Noordegraaf A, Beghetti M, Ghofrani A, Gomez Sanchez MA, Hansmann G, Klepetko W, Lancellotti P, Matucci M, McDonagh T, Pierard LA, Trindade PT, Zompatori M, Hoeper M. 2015 ESC/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension. *Eur Respir J* 2015;46:903–75.
- Chin KM, Rubin LJ. Pulmonary arterial hypertension. *J Am Coll Cardiol* 2008;51:1527–38.
- Humbert M, Sitbon O, Chaouat A, Bertocchi M, Habib G, Gressin V, Yaici A, Weitzenblum E, Cordier J-F, Chabot F, Dromer C, Pison C, Reynaud-Gaubert M, Haloun A, Laurent M, Hachulla E, Simonneau G. Pulmonary arterial hypertension in France. *Am J Respir Crit Care Med* 2006;173:1023–30.
- Dresdale DT, Schultz M, Michtom RJ. Primary pulmonary hypertension. I. Clinical and hemodynamic study. *Am J Med* 1951;11:686–705.
- Lane KB, Machado RD, Paucillo MW, Thomson JR, Phillips JA, Loyd JE, Nichols WC, Trembath RC. International PPH Consortium. Heterozygous germline mutations in *BMPR2*, encoding a TGF-beta receptor, cause familial primary pulmonary hypertension. *Nat Genet* 2000;26:81–4.
- White RJ, Morrell NW. Understanding the low penetrance of bone morphogenetic protein receptor 2 gene mutations. *Circulation* 2012;126:1818–20.
- Austin ED, Loyd JE. The genetics of pulmonary arterial hypertension. *Circ Res* 2014;115:189–202.
- Girerd B, Montani D, Coulet F, Sztrymf B, Yaici A, Jais X, Tregouët D, Reis A, Drouin-Garraud V, Fraisse A, Sitbon O, O'Callaghan DS, Simonneau G, Soubrier F, Humbert M. Clinical outcomes of pulmonary arterial hypertension in patients carrying an ACVRL1 (ALK1) mutation. *Am J Respir Crit Care Med* 2010;181:851–61.
- Pfarr N, Fischer C, Ehlen N, Becker-Grünig T, López-González V, Gorenflo M, Hager A, Hinderhofer K, Miera O, Nagel C, Schranz D, Grünig E. Hemodynamic and genetic analysis in children with idiopathic, heritable, and congenital heart disease associated pulmonary arterial hypertension. *Respir Res* 2013;14:3.
- Shintani M, Yagi H, Nakayama T, Saji T, Matsuoka R. A new nonsense mutation of *SMAD8* associated with pulmonary arterial hypertension. *J Med Genet* 2009;46:331–7.
- Austin ED, Ma L, LeDuc C, Berman Rosenzweig E, Borczuk A, Phillips JA, Palomero T, Sumazin P, Kim HR, Talati MH, West J, Loyd JE, Chung WK. Whole exome sequencing to identify a novel gene (caveolin-1) associated with human pulmonary arterial hypertension. *Circ Cardiovasc Genet* 2012;5:336–43.
- Ma L, Roman-Campos D, Austin ED, Eyries M, Sampson KS, Soubrier F, Germain M, Tréguët DA, Borczuk A, Rosenzweig EB, Girerd B, Montani D, Humbert M, Loyd JE, Kass RS, Chung WK. A novel channelopathy in pulmonary arterial hypertension. *N Engl J Med* 2013;369:351–61.
- Newman JH, Trembath RC, Morse JA, Grunig E, Loyd JE, Adnot S, Cocco F, Ventura C, Phillips JA, Knowles JA, Janssens B, Eickelberg O, Eddahibi S, Herve P, Nichols WC, Elliott G. Genetic basis of pulmonary arterial hypertension: current understanding and future directions. *J Am Coll Cardiol* 2004;43:533–9.
- Cooper DN, Krawczak M, Polychronakos C, Tyler-Smith C, Kehrer-Sawatzki H. Where genotype is not predictive of phenotype: towards an understanding of the molecular basis of reduced penetrance in human inherited disease. *Hum Genet* 2013;132:1077–130.
- West J, Cogan J, Geraci M, Robinson L, Newman J, Phillips JA, Lane K, Meyrick B, Loyd J. Gene expression in *BMPR2* mutation carriers with and without evidence of pulmonary arterial hypertension suggests pathways relevant to disease penetrance. *BMJ Med Genomics* 2008;1:45.
- Hamid R, Cogan JD, Hedges LK, Austin E, Phillips JA, Newman JH, Loyd JE. Penetrance of pulmonary arterial hypertension is modulated by the expression of normal *BMPR2* allele. *Hum Mutat* 2009;30:649–54.
- Germain M, Eyries M, Montani D, Poirier O, Girerd B, Dorfmueller P, Coulet F, Nadaud S, Maugendre S, Guignabert C, Carpentier W, Vonk-Noordegraaf A, Lévy M, Chaouat A, Lambert JC, Bertrand M, Dupuy AM, Letenneur L, Lathrop M, Amouyel P, de Ravel TJ, Delcroix M, Austin ED, Robbins IM, Hemmes AR, Loyd JE, Berman-Rosenzweig E, Barr J, Chung WK, Simonneau G, Tréguët DA, Humbert M, Soubrier F. Genome-wide association analysis identifies a susceptibility locus for pulmonary arterial hypertension. *Nat Genet* 2013;45:518–21.
- Eichstaedt CA, Song J, Benjamin N, Harutyunova S, Fischer C, Grünig E, Hinderhofer K. EIF2AK4 mutation as "second hit" in hereditary pulmonary arterial hypertension. *Respir Res* 2016;17:141.
- Viales RR, Eichstaedt CA, Ehlen N, Fischer C, Lichtblau M, Grünig E, Hinderhofer K. Mutation in *BMPR2* promoter: a 'second hit' for manifestation of pulmonary arterial hypertension? *PLoS One* 2015;10:e0133042.
- Aldred MA, Comhair SA, Varela-García M, Asosingh K, Xu W, Noon GP, Thistlethwaite PA, Tudor RM, Erzurum SC, Geraci MW, Coldren CD. Somatic chromosome abnormalities in the lungs of patients with pulmonary arterial hypertension. *Am J Respir Crit Care Med* 2010;182:1153–60.
- Austin ED, Cogan JD, West JD, Hedges LK, Hamid R, Dawson EP, Wheeler LA, Parl FF, Loyd JE, Phillips JA. Alterations in oestrogen metabolism: implications for higher penetrance of familial pulmonary arterial hypertension in females. *Eur Respir J* 2009;34:1093–9.

- 22 Portillo K, Santos S, Madrigal I, Blanco I, Paré C, Borderías L, Peinado VI, Roca J, Milà M, Barberà JA. Study of the BMP2 gene in patients with pulmonary arterial hypertension. *Archivos de Bronconeumología* 2010;46:129–34.
- 23 Deng Z, Morse JH, Slager SL, Cuervo N, Moore KJ, Venetos G, Kalachikov S, Cayanis E, Fischer SG, Barst RJ, Hodge SE, Knowles JA. Familial primary pulmonary hypertension (gene PPH1) is caused by mutations in the bone morphogenetic protein receptor-II gene. *Am J Hum Genet* 2000;67:737–44.
- 24 Rudarakanchana N, Flanagan JA, Chen H, Upton PD, Machado R, Patel D, Trembath RC, Morrell NW. Functional analysis of bone morphogenetic protein type II receptor mutations underlying primary pulmonary hypertension. *Hum Mol Genet* 2002;11:1517–25.
- 25 Miller SA, Dykes DD, Polesky HF. A simple salting out procedure for extracting DNA from human nucleated cells. *Nucleic Acids Res* 1988;16:1215.
- 26 Lek M, Karczewski KJ, Minikel EV, Samocha KE, Banks E, Fennell T, O'Donnell-Luria AH, Ware JS, Hill AJ, Cummings BB, Tukiainen T, Birnbaum DP, Kosmicki JA, Duncan LE, Estrada K, Zhao F, Zou J, Pierce-Hoffman E, Berghout J, Cooper DN, DeLafont N, DePristo M, Do R, Flannick J, Fromer M, Gauthier L, Goldstein J, Gupta N, Howrigan D, Kiezun A, Kurki MI, Moonshine AL, Natarajan P, Orozco L, Peloso GM, Poplin R, Rivas MA, Ruano-Rubio V, Rose SA, Ruderfer DM, Shakir K, Stenson PD, Stevens C, Thomas BP, Tiao G, Tusie-Luna MT, Weisburd B, Won HH, Yu D, Altshuler DM, Ardisson D, Boehnke M, Danesh J, Donnelly S, Elosua R, Florez JC, Gabriel SB, Getz G, Glatt SJ, Hultman CM, Kathiresan S, Laakso M, McCarroll S, McCarthy MI, McGovern D, McPherson R, Neale BM, Palotie A, Purcell SM, Saleheen D, Scharf JM, Sklar P, Sullivan PF, Tuomilehto J, Tsuang MT, Watkins HC, Wilson JG, Daly MJ, MacArthur DG. Exome Aggregation Consortium. Analysis of protein-coding genetic variation in 60,706 humans. *Nature* 2016;536:285–91.
- 27 Auton A, Brooks LD, Durbin RM, Garrison EP, Kang HM, Korbel JO, Marchini JL, McCarthy S, McVean GA, Abecasis GR; 1000 Genomes Project Consortium. A global reference for human genetic variation. *Nature* 2015;526:68–74.
- 28 Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MA, Bender D, Maller J, Sklar P, de Bakker PI, Daly MJ, Sham PC. PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet* 2007;81:559–75.
- 29 Matise TC, Chen F, Chen W, De La Vega FM, Hansen M, He C, Hyland FC, Kennedy GC, Kong X, Murray SS, Ziegler JS, Stewart WC, Buysse S. A second-generation combined linkage physical map of the human genome. *Genome Res* 2007;17:1783–6.
- 30 Lange K, Papp JC, Sinsheimer JS, Sripracha R, Zhou H, Sobel EM. Mendel: the Swiss army knife of genetic analysis programs. *Bioinformatics* 2013;29:1568–70.
- 31 Gertz EM, Hiekkalinna T, Digabel SL, Audet C, Terwilliger JD, Schäffer AA. PSEUDOMARKER 2.0: efficient computation of likelihoods using NOMAD. *BMC Bioinformatics* 2014;15:47.
- 32 Silberstein M, Weissbrod O, Otten L, Tzemach A, Anisania A, Shtark O, Tuberg D, Galfrin E, Gannon I, Shalata A, Borochowitz ZU, Dechter R, Thompson E, Geiger D. A system for exact and approximate genetic linkage analysis of SNP data in large pedigrees. *Bioinformatics* 2013;29:197–205.
- 33 Abecasis GR, Cherny SS, Cookson WO, Cardon LR. Merlin—rapid analysis of dense genetic maps using sparse gene flow trees. *Nat Genet* 2002;30:97–101.
- 34 Wijsman EM, Rothstein JH, Thompson EA. Multipoint linkage analysis with many multiallelic or dense diallelic markers: Markov chain-Monte Carlo provides practical approaches for genome scans on general pedigrees. *Am J Hum Genet* 2006;79:846–58.
- 35 MacArthur J, Bowler E, Cerezo M, Gil L, Hall P, Hastings E, Junkins H, McMahon A, Milano A, Morales J, Pendlington ZM, Welter D, Burdett T, Hindorf L, Flicek P, Cunningham F, Parkinson H. The new NHGRI-EBI Catalog of published genome-wide association studies (GWAS Catalog). *Nucleic Acids Res* 2017;45:D896–901.
- 36 Kato N, Takeuchi F, Tabara Y, Kelly TN, Go MJ, Sim X, Tay WT, Chen CH, Zhang Y, Yamamoto K, Katsuya T, Yokota M, Kim YJ, Ong RT, Nabika T, Gu D, Chang LC, Kokubo Y, Huang W, Ohnaka K, Yamori Y, Nakashima E, Jaquish CE, Lee JY, Seielstad M, Isono M, Hixson JE, Chen YT, Miki T, Zhou X, Sugiyama T, Jeon JP, Liu JJ, Takayanagi R, Kim SS, Aung T, Sung YJ, Zhang X, Wong TY, Han BG, Kobayashi S, Ogihara T, Zhu D, Iwai N, Wu JY, Teo YY, Tai ES, Cho YS, He J. Meta-analysis of genome-wide association studies identifies common variants associated with blood pressure variation in east Asians. *Nat Genet* 2011;43:531–8.
- 37 Kato N, Loh M, Takeuchi F, Verweij N, Wang X, Zhang W, Kelly TN, Saleheen D, Lehne B, Leach IM, Drong AW, Abbott J, Wahl S, Tan ST, Scott WR, Campanella G, Chadeau-Hyam M, Afzal U, Ahluwalia TS, Bonder MJ, Chen P, Dehghan A, Edwards TL, Esko T, Go MJ, Harris SE, Hartiaia J, Kasela S, Kasturiratne A, Khor CC, Kleber ME, Li H, Yu Mok Z, Nakatochi M, Sapari NS, Saxena R, Stewart AFR, Stolk L, Tabara Y, Teh AL, Wu Y, Wu JY, Zhang Y, Aits I, Da Silva Couto Alves A, Das S, Dorajoo R, Hopewell JC, Kim YK, Koivula RW, Luan J, Lyttikainen LP, Nguyen QN, Pereira MA, Postmus I, Raitakari OT, Scannell Bryan M, Scott RA, Sorice R, Tragante V, Traglia M, White J, Yamamoto K, Zhang Y, Adair LS, Ahmed A, Akiyama K, Asif R, Aung T, Barroso I, Bjorntjes A, Braun TR, Cai H, Chang LC, Chen CH, Cheng CY, Chong YS, Collins R, Courtney R, Davies G, Delgado G, Do LD, Doevendans PA, Gansevoort RT, Gao YT, Grammer TB, Grarup N, Grewal J, Gu D, Wander GS, Hartikainen AL, Hazen SL, He J, Heng CK, Hixson JE, Hofman A, Hsu C, Huang W, Husemoen LLN, Hwang JY, Ichihara S, Igase M, Isono M, Justesen JM, Katsuya T, Kibria MG, Kim YJ, Kishimoto M, Koh WP, Kohara K, Kumari M, Kwek K, Lee NR, Lee J, Liao J, Lieb W, Liewald DCM, Matsubara T, Matsushita Y, Meitinger T, Mihailov E, Milani L, Mills R, Mononen N, Müller-Nurasyid M, Nabika T, Nakashima E, Ng HK, Nikus K, Nutile T, Ohkubo T, Ohnaka K, Parish S, Paternoster L, Peng H, Peters A, Pham ST, Piniidiapathirage MJ, Rahman M, Rakugi H, Rolandsson O, Ann Rozario M, Ruggiero D, Sala CF, Sarju R, Shimokawa K, Snieder H, Sparsa T, Spiering WW, Starr JM, Stott DJ, Stram DO, Sugiyama T, Szymczak S, Tang WHW, Tong L, Trompet S, Turjanmaa V, Ueshima H, Uitterlinden AG, Umemura S, Vaarasmaki M, van Dam RM, van Gilst WH, van Veldhuisen DJ, Viikari JS, Waldenberger M, Wang Y, Wang A, Wilson R, Wong TY, Xiang YB, Yamaguchi S, Ye X, Young RD, Young TL, Yuan JM, Zhou X, Asselbergs FW, Ciullo M, Clarke R, Deloukas P, Franke A, Franks PW, Franks S, Friedlander Y, Gross MD, Guo Z, Hansen T, Jarvelin MR, Jørgensen T, Jukema JW, Kähönen M, Kajio H, Kimviki M, Lee JY, Lehtimäki T, Linneberg A, Miki T, Pedersen O, Samani NJ, Sørensen TIA, Takayanagi R, Toniolo D, Ahsan H, Allayee H, Kuan YT, Danesh J, Deary IJ, Franco OH, Franke L, Heijman BT, Holbrook JD, Isaacs A, Kim BJ, Lin X, Liu J, März W, Metspalu A, Mohlke KL, Sanghera DK, Shu XO, van Meurs JBJ, Vithana E, Wickremasinghe AR, Wijmenga C, Wolfenbutter BHW, Yokota M, Zheng W, Zhu D, Vineis P, Kyrtopoulos SA, Kleijmans JC, McCarthy MI, Soong R, Gieger C, Scott J, Teo YY, He J, Elliott P, Tai ES, van der Harst P, Kooner JS, Chambers JC. BIOS-consortium/CARDIo GRAMplus/CdLifeLines Cohort Study/InterAct Consortium. Trans-ancestry genome-wide association study identifies 12 genetic loci influencing blood pressure and implicates a role for DNA methylation. *Nat Genet* 2015;47:1282–93.
- 38 Simino J, Shi G, Bis JC, Chasman DI, Ehret GB, Gu X, Guo X, Hwang SJ, Sijbrands E, Smith AV, Verwoert GC, Bragg-Gresham JL, Cady G, Chen P, Cheng CY, Corre T, de Boer RA, Goel A, Johnson T, Khor CC, Lluís-Ganella C, Luan J, Lyttikainen LP, Nolte IM, Sim X, Söber S, van der Most PJ, Verweij N, Zhao JH, Amin N, Boerwinkle E, Bouchard C, Dehghan A, Eiriksdottir G, Elosua R, Franco OH, Gieger C, Harris TB, Herberg S, Hofman A, James AL, Johnson AD, Kähönen M, Khaw KT, Kutalik Z, Larson MG, Launer LJ, Li G, Liu J, Liu K, Morrison AC, Navis G, Ong RT, Papanicolaou GJ, Penninx BW, Psaty BM, Raffel LJ, Raitakari OT, Rice K, Rivadeneira F, Rose LM, Sanna S, Scott RA, Siscovick DS, Stolk RP, Uitterlinden AG, Vaidya D, van der Klauw MM, Vasani RS, Vithana EN, Völker U, Völzke H, Watkins H, Young TL, Aung T, Bochud M, Farrall M, Hartman CA, Laan M, Lakatta EG, Lehtimäki T, Loos RJ, Lucas G, Meneton P, Palmer LJ, Rettig R, Snieder H, Tai ES, Teo YY, van der Harst P, Wareham NJ, Wijmenga C, Wong TY, Fornage M, Gudnason V, Levy D, Palmas W, Ridker PM, Rotter JJ, van Duijn CM, Witteman JC, Chakravarti A, Rao DC. LifeLines Cohort Study. Gene-age interactions in blood pressure regulation: a large-scale investigation with the CHARGE, Global BPgen, and ICBP Consortia. *Am J Hum Genet* 2014;95:24–38.
- 39 Wain LV, Vaez A, Jansen R, Joehanes R, van der Most PJ, Erzurumluoglu AM, O'Reilly PF, Cabrera CP, Warren HR, Rose LM, Verwoert GC, Hottenga JJ, Strawbridge RJ, Esko T, Arking DE, Hwang SJ, Guo X, Kutalik Z, Trompet S, Shrine N, Teumer A, Ried JS, Bis JC, Smith AV, Amin N, Nolte IM, Lyttikainen LP, Mahajan A, Wareham NJ, Wijmenga C, Joshi PK, Kristiansson K, Traglia M, Havulinna AS, Goel A, Nalls MA, Söber S, Vuckovic D, Luan J, Del Greco M F, Ayers KL, Marrugat J, Ruggiero D, Lopez LM, Niiranen T, Enroth S, Jackson AU, Nelson CP, Huffman JE, Zhang W, Marten J, Gandin I, Harris SE, Zemunik T, Lu Y, Evangelou E, Shah N, de Borst MH, Mangino M, Prins BP, Campbell A, Li-Gao R, Chauhan G, Oldmeadow C, Abecasis G, Abedi M, Barbieri CM, Barnes MR, Batini C, Beilby J, Blake T, Boehnke M, Bottinger EP, Braund PS, Brown M, Brumat M, Campbell H, Chambers JC, Cocca M, Collins F, Connell J, Cordell HJ, Damman JJ, Davies G, de Geus EJ, de Mutser R, Deelen J, Demirkan K, Doney ASF, Dör M, Farrall M, Ferreira T, Fränberg M, Gao H, Giedraitis V, Gieger C, Giulianini F, Gow AJ, Hamsten A, Harris TB, Hofman A, Holliday EG, Hui J, Jarvelin MR, Johansson Å, Johnson AD, Jousilahti P, Jula A, Kähönen M, Kathiresan S, Khaw KT, Kolcic I, Kosken S, Langenberg C, Larson M, Launer LJ, Lehne B, Liewald DCM, Lin L, Lind L, Mach F, Mamasoula C, Menni C, Mifsud B, Milanesechi Y, Morgan A, Morris AD, Morrison AC, Munson PJ, Nandakumar P, Nguyen QT, Nutile T, Oldehinkel AJ, Oostra BA, Orger E, Padmanabhan S, Palotie A, Paré G, Pattie A, Penninx B, Poulter N, Promtstaller PP, Raitakari OT, Ren M, Rice K, Ridker PM, Riese H, Ripatti S, Robino A, Rotter JJ, Rudan I, Saba Y, Saint Pierre A, Sala CF, Sarin AP, Schmidt R, Scott R, Seelen MA, Shields DC, Siscovick D, Sorice R, Stanton A, Stott DJ, Sundström J, Swertz M, Taylor KD, Thom S, Tzoulaki I, Tzourio C, Uitterlinden AG, Völker U, Vollenweider P, Wild S, Willemssen G, Wright AF, Yao J, Thériault S, Conen D, Attia J, Sever P, DeBette S, Mook-Kanamori DO, Zeggini E, Spector TD, van der Harst P, Palmer CNA, Vergnaud AC, Loos RJF, Polasek O, Starr JM, Grotto G, Hayward C, Kooner JS, Lindgren CM, Vitart V, Samani NJ, Tuomilehto J, Gyllenstein U, Knekt P, Deary IJ, Ciullo M, Elosua R, Keavney BD, Hicks AA, Scott RA, Gasparini P, Laan M, Liu Y, Watkins H, Hartman CA, Salomaa V, Toniolo D, Perola M, Wilson JF, Schmidt H, Zhao JH, Lehtimäki T, van Duijn CM, Gudnason V, Psaty BM, Peters A, Rettig R, James A, Jukema JW, Strachan DP, Palmas W, Metspalu A, Ingelsson E, Boomsma DI, Franco OH, Bochud M, Newton-Cheh C, Munroe PB, Elliott P, Chasman DI, Chakravarti A, Knight J, Morris AP, Levy D, Tobin MD, Snieder H, Caulfield MJ, Ehret GB. Novel blood pressure locus and gene discovery using genome-wide association study and expression data sets from blood and the kidney. *Hypertension* 2017.
- 40 Wain LV, Verwoert GC, O'Reilly PF, Shi G, Johnson T, Johnson AD, Bochud M, Rice KM, Henneman P, Smith AV, Ehret GB, Amin N, Larson MG, Mooser V, Hadley D, Dör M, Bis JC, Aspelund T, Esko T, Janssens AC, Zhao JH, Heath S, Laan M, Fu J, Pistis G, Luan J, Arora P, Lucas G, Pirastu N, Pichler I, Jackson AU, Webster RJ, Zhang F, Peden JF, Schmidt H, Tanaka T, Campbell H, Igl W, Milanesechi Y, Hottenga JJ, Vitart V, Chasman DI, Trompet S, Bragg-Gresham JL, Alizadeh BZ, Chambers JC, Guo X, Lehtimäki T, Kühnel B, Lopez LM, Polašek O, Boban M, Nelson CP, Morrison AC, Pihur V, Ganesh SK, Hofman A, Kundu S, Mattace-Raso FU, Rivadeneira F, Sijbrands EJ, Uitterlinden

- AG, Hwang SJ, Vasan RS, Wang TJ, Bergmann S, Vollenweider P, Waeber G, Laitinen J, Pouta A, Zitting P, McArdle WL, Kroemer HK, Völker U, Völzke H, Glazer NL, Taylor KD, Harris TB, Alavere H, Haller T, Keis A, Tammesoo ML, Aulchenko Y, Barroso I, Khaw KT, Galan P, Hercberg S, Lathrop M, Eyheramendy S, Org E, Söber S, Lu X, Nolte IM, Penninx BW, Corre T, Masciullo C, Sala C, Groop L, Voight BF, Melander O, O'Donnell CJ, Salomaa V, d'Adamo AP, Fabretto A, Faletra F, Ulivi S, Del Greco F, Facheris M, Collins FS, Bergman RN, Beilby JP, Hung J, Musk AW, Mangino M, Shin SY, Soranzo N, Watkins H, Goel A, Hamsten A, Gider P, Loitfelder M, Zeginigg M, Hernandez D, Najjar SS, Navarro P, Wild SH, Corsi AM, Singleton A, de Geus EJ, Willemsen G, Parker AN, Rose LM, Buckley B, Stott D, Orru M, Uda M, van der Klauw MM, Zhang W, Li X, Scott J, Chen YD, Burke GL, Kähönen M, Viikari J, Döring A, Meitinger T, Davies G, Starr JM, Emilsson V, Plump A, Lindeman JH, Hoen PA, König IR, Felix JF, Clarke R, Hopewell JC, Ongen H, Breteler M, Debetto S, Destefano AL, Fornage M, Mitchell GF, Smith NL, Holm H, Stefansson K, Thorleifsson G, Thorsteinsdottir U, Samani NJ, Preuss M, Rudan I, Hayward C, Deary IJ, Wichmann HE, Raitakari OT, Palmas W, Kooner JS, Stolk RP, Jukema JW, Wright AF, Boomsma DI, Bandinelli S, Gyllenstein UB, Wilson JF, Ferrucci L, Schmidt R, Farrall M, Spector TD, Palmer LJ, Tuomilehto J, Pfeufer A, Gasparini P, Siscovick D, Altshuler D, Loos RJ, Toniolo D, Snieder H, Gieger C, Meneton P, Wareham NJ, Oostra BA, Metspalu A, Launer L, Rettig R, Strachan DP, Beckmann JS, Witteman JC, Erdmann J, van Dijk KW, Boerwinkle E, Boehnke M, Ridker PM, Jarvelin MR, Chakravarti A, Abecasis GR, Gudnason V, Newton-Cheh C, Levy D, Munroe PB, Psaty BM, Caulfield MJ, Rao DC, Tobin MD, Elliott P, van Duijn CM. LifeLines Cohort StudyEchoGen consortiumAortaGen ConsortiumCHARGE Consortium Heart Failure Working GroupKidneyGen consortiumCKDGen consortiumCardiogenics consortiumCardioGram. Genome-wide association study identifies six new loci influencing pulse pressure and mean arterial pressure. *Nat Genet* 2011;43:1005–11.
- 41 Yucelsoy B, Kaufman KM, Lummus ZL, Weirauch MT, Zhang G, Cartier A, Boulet LP, Sastre J, Quirce S, Tarlo SM, Cruz MJ, Munoz X, Harley JB, Bernstein DI. Genome-wide association study identifies novel loci associated with diisocyanate-induced occupational asthma. *Toxicol Sci* 2015;146:192–201.
- 42 Higgins M, Province M, Heiss G, Eckfeldt J, Ellison RC, Folsom AR, Rao DC, Sprafka JM, Williams R. NHLBI Family Heart Study: objectives and design. *Am J Epidemiol* 1996;143:1219–28.
- 43 GTEx Consortium. Human genomics. The Genotype-Tissue Expression (GTEx) pilot analysis: multitissue gene regulation in humans. *Science* 2015;348:648–60.
- 44 Schmitt AD, Hu M, Jung I, Xu Z, Qiu Y, Tan CL, Li Y, Lin S, Lin Y, Barr CL, Ren B. A compendium of chromatin contact maps reveals spatially active regions in the human genome. *Cell Rep* 2016;17:2042–59.
- 45 Ward LD, Kellis M. HaploReg: a resource for exploring chromatin states, conservation, and regulatory motif alterations within sets of genetically linked variants. *Nucleic Acids Res* 2012;40:D930–4.
- 46 Moyerbrailean GA, Kalita CA, Harvey CT, Wen X, Luca F, Pique-Regi R. Which genetics variants in DNase-Seq footprints are more likely to alter binding? *PLoS Genet* 2016;12:e1005875.
- 47 Zhao YD, Yun HZH, Peng J, Yin L, Chu L, Wu L, Michalek R, Liu M, Keshavjee S, Waddell T, Granton J, de Perrot M. De novo synthesis of bile acids in pulmonary arterial hypertension lung. *Metabolomics* 2014;10:1169–75.
- 48 Mukherjee S, Diaz Valencia JD, Stewman S, Metz J, Monnier S, Rath U, Asenjo AB, Charafeddine RA, Sosa HJ, Ross JL, Ma A, Sharp DJ. Human Fidgetin is a microtubule severing the enzyme and minus-end depolymerase that regulates mitosis. *Cell Cycle* 2012;11:2359–66.
- 49 Wang D, Wang F, Shi KH, Tao H, Li Y, Zhao R, Lu H, Duan W, Qiao B, Zhao SM, Wang H, Zhao JY. Lower circulating folate induced by a Fidgetin intronic variant is associated with reduced congenital heart disease susceptibility. *Circulation* 2017;135:1733–48.
- 50 Devoto M, Specchia C, Laudenslager M, Longo L, Hakonarson H, Maris J, Mossé Y. Genome-wide linkage analysis to identify genetic modifiers of ALK mutation penetrance in familial neuroblastoma. *Hum Hered* 2011;71:135–9.