

Supplementary File 2. Data from case reports of individuals with pathogenic variants in autism-associated genes

Autism gene	ID cases	Autism cases	Autism cases with comorbid ID	Autism cases without ID	Variant types in autism/ID cases	Ref.
<i>ADNP</i>	11/11	7/11	7/7	0/7	Unknown	[1]
	Mild - 9/73 Moderate - 26/73 Severe - 38/73 TOTAL - 73/73	64/73	64/64	0/64	Unknown	[2]
	1/1 - Severe	1/1	1/1	0/1	Nonsense	[3]
	36/36	29/36	29/29	0/29	Missense - 1 Nonsense - 14 Frameshift - 13 Deletion - 1	[4]
	13/13	13/13	13/13	0/13	Unknown	[5]
<i>ANK2</i>	1/1 - Moderate (IQ=40-60)	1/1	1/1	0/1	Frameshift	[6]
<i>ANKRD11</i>	Moderate - 6/7 Mild - 1/7 TOTAL - 7/7	7/7	7/7	0/7	Nonsense - 1 Frameshift - 3 Deletions - 3	[7]
	1/1	1/1	1/1	0/1	Frameshift	[8]
	2/2	1/2	1/1	0/1	Nonsense	[9]
<i>ARID1B</i>	127/143	77/143	77/77	0/77	LOF - 118 Deletions - 18 Splice site - 7	[10]
	1/1 - Severe	1/1	1/1	0/1	Frameshift	[11]
	8/8 - Moderate/Severe	1/8	1/1	0/1	Nonsense	[12]
	1/1	1/1	1/1	0/1	Frameshift	[13]

Autism gene	ID cases	Autism cases	Autism cases with comorbid ID	Autism cases without ID	Variant types in autism/ID cases	Ref.
<i>ASH1L</i>	6/13 - Mild 5/13 - Severe 2/13 - Yes TOTAL - 13/13	3/13	3/3	0/3	Nonsense - 2 Frameshift - 1	[14]
	1/1 - IQ<30	1/1	1/1	0/1	Missense	[15]
<i>ASXL3</i>	1/1	1/1	1/1	0/1	Nonsense	[16]
	3/12 - Moderate 1/12 - Moderate-Severe 6/12 - Severe 1/12 - Profound TOTAL - 11/12	9/12	8/9	1/9	Nonsense - 6 Frameshift - 2	[17]
	1/1 - Severe	1/1	1/1	0/1	Frameshift	[18]
	1/1 - Profound	1/1	1/1	0/1	Frameshift	[19]
	2/2 - Severe	2/2	2/2	0/2	Nonsense - 2	[20]
	1/1	1/1	1/1	0/1	Biallelic Missense	[21]
	1/1	1/1	1/1	0/1	Frameshift	[22]
<i>BCL11A</i>	1/9 - Mild-Moderate 1/9 - Moderate-Severe 6/9 - Moderate 1/9 - Severe TOTAL - 9/9	2/9	2/2	0/2	Missense - 1 Frameshift - 1	[23]
	2/7	2/7	1/2	1/2	Missense	[24]
<i>CHD2</i>	1/1	1/1	1/1	0/1	Frameshift	[25]
	2/2	2/2	2/2	0/2	Nonsense - 1 Splicing - 2	[26]

Autism gene	ID cases	Autism cases	Autism cases with comorbid ID	Autism cases without ID	Variant types in autism/ID cases	Ref.
<i>CHD8</i>	1/10 - Mild 3/10 - Moderate 1/10 - Moderate-Severe 2/10 - Severe TOTAL - 7/10	7/10	6/7	1/7	Nonsense - 4 Splicing - 1 Frameshift - 1	[27]
	7/10	9/10	6/9	3/9	Missense - 2 Nonsense - 4	[28]
	11/11	6/11	6/6	0/6	Unknown	[1]
	1/2 - Severe 1/2 - IQ=33 TOTAL - 2/2	2/2	2/2	0/2	Frameshift - 2	[15]
	11/17	14/17	9/14	5/14	Deletion - 1 Nonsense - 1 Splicing - 1 Frameshift - 3	[29]
	9/15	13/15	8/13	5/13	Nonsense - 4 Splicing - 1 Frameshift - 3	[30]
	1/10 - Below average IQ 1/10 - Mild 3/10 - Low IQ 4/10 - Extremely low IQ TOTAL - 9/10	10/10	9/10	1/10	Deletions - 1 Nonsense - 3 Splicing - 1 Frameshift - 4	[31]
<i>CUL3</i>	1/1 - Mild	1/1	1/1	0/1	Missense	[32]
<i>DDX3X</i>	69/69	27/69	27/27	0/27	Unknown	[33]
	28/28	6/28	6/6	0/6	Unknown	[34]

Autism gene	ID cases	Autism cases	Autism cases with comorbid ID	Autism cases without ID	Variant types in autism/ID cases	Ref.
<i>DYRK1A</i>	6/6	6/6	6/6	0/6	Nonsense - 2 Frameshift - 4	[35]
	1/20	3/20	1/3	2/3	Nonsense	[36]
<i>KMT2A</i>	5/6	5/6	4/5	1/5	Missense - 1 Nonsense - 1 Frameshift - 2	[37]
	0/2	1/2	0/1	1/1	Splicing - 1	[38]
	18/33 - Mild 11/33 - Moderate 4/33 - Severe TOTAL - 33/33	2/33	2/2	0/2	Missense - 2	[39]
	38/58	4/58	4/4	0/4	Unknown	[40]
<i>MECP2</i>	1/1	1/1	1/1	0/1	Nonsense	[41]
<i>MYT1L</i>	1/1 - IQ=60	1/1	1/1	0/1	Nonsense	[15]
<i>NRXN1</i>	11/11 - Severe	1/11	1/1	0/1	Bi-allelic/deletion	[42]
	8/21	12/21	2/12	10/12	Deletions - 2	[43]
	25/27	20/27	19/20	1/20	Deletions - 19	[44]
	1/1	1/1	1/1	0/1	Deletion	[45]
<i>POGZ</i>	9/10	5/10	5/5	0/5	Missense - 1 Nonsense - 3 Duplication - 1	[46]
	1/1	1/1	1/1	0/1	Missense	[47]
	1/1	1/1	1/1	0/1	LOF	[48]
	1/1 - Severe	1/1	1/1	0/1	Frameshift	[49]
	24/25	17/23	17/17	0/17	Intronic - 2 Nonsense - 7 Frameshift - 8	[50]

Autism gene	ID cases	Autism cases	Autism cases with comorbid ID	Autism cases without ID	Variant types in autism/ID cases	Ref.
POGZ	5/5	2/5	2/2	0/2	Nonsense - 1 Frameshift - 1	[51]
	6/6	1/7	1/1	0/1	Nonsense	[52]
	1/1 - Severe	1/1	1/1	0/1	Missense	[53]
SCN2A	1/1	1/1	1/1	0/1	Deletion	[54]
	1/1	1/1	1/1	0/1	Missense	[55]
	1/1 - Severe	1/1	1/1	0/1	Missense	[56]
	1/1	1/1	1/1	0/1	Splicing	[57]
	15/33	5/33	3/5	2/5	Missense - 2 Nonsense - 1	[58]
SETD5	8/9	2/9	2/2	0/2	Intronic - 1 Nonsense - 1	[59]
	7/7	5/7	5/5	0/5	Unknown	[60]
SHANK3	29/30	22/30	21/22	1/22	Missense - 1 Splicing - 2 Nonsense - 4 Frameshift - 14	[61]
	0/1	1/1	0/1	1/1	Frameshift	[62]
	1/1	1/1	1/1	0/1	Frameshift	[63]
	5/5	2/5	2/2	0/2	Unknown	[64]
SYNGAP1	1/1 - Moderate	1/1	1/1	0/1	Frameshift	[65]
	1/8 - Moderate 6/8 - Moderate/Severe 1/8 - Severe TOTAL - 8/8	3/8	3/3	0/3	Missense - 2 Frameshift - 1	[66]
	1/1	1/1	1/1	0/1	Splicing	[67]
	1/1 - Mild	1/1	1/1	0/1	Missense	[68]
	1/1 - Severe	1/1	1/1	0/1	Splicing	[69]

Autism gene	ID cases	Autism cases	Autism cases with comorbid ID	Autism cases without ID	Variant types in autism/ID cases	Ref.
<i>SYNGAP1</i>	1/11 - Mild 6/11 - Moderate 4/11 - Moderate/Severe TOTAL - 11/11	4/11	4/4	0/4	Missense - 2 Splicing - 1 Frameshift - 1	[70]
<i>TCF20</i>	16/16	9/16	9/9	0/9	Missense - 1 Nonsense - 1 Deletion - 3 Frameshift - 4	[71]
	26/26	18/26	18/18	0/18	Unknown	[72]
	1/5 - borderline (IQ=79) 2/5 - Mild 1/5 - Mild-Moderate 1/5 - Moderate TOTAL - 5/6	5/6	4/5	1/5	Rearrangement - 2 Frameshift - 2	[73]

References

- 1 Arnett AB, Rhoads CL, Hoekzema K, *et al.* The autism spectrum phenotype in ADNP syndrome. *Autism Res* 2018;**11**:1300–10. doi:10.1002/aur.1980
- 2 Van Dijck A, Vulto-van Silfhout AT, Cappuyns E, *et al.* Clinical Presentation of a Complex Neurodevelopmental Disorder Caused by Mutations in ADNP. *Biol Psychiatry* 2019;**85**:287–97. doi:10.1016/j.biopsych.2018.02.1173
- 3 Alkhunaizi E, Walkiewicz MA, Chitayat D. Mutation in the ADNP gene associated with Noonan syndrome features. *Clin Dysmorphol* 2018;**27**:53–7. doi:10.1097/MCD.0000000000000215
- 4 Gozes I, Van Dijck A, Hacoen-Kleiman G, *et al.* Premature primary tooth eruption in cognitive/motor-delayed ADNP-mutated children. *Transl Psychiatry* 2017;**7**. doi:10.1038/tp.2017.27
- 5 Krajewska-Walasek M, Jurkiewicz D, Piekutowska-Abramczuk D, *et al.* Additional data on the clinical phenotype of Helsmoortel-Van der Aa syndrome associated with a novel truncating mutation in ADNP gene. *Am J Med Genet Part A* 2016;**170**:1647–50. doi:10.1002/ajmg.a.37641
- 6 Bowling KM, Thompson ML, Amaral MD, *et al.* Genomic diagnosis for children with intellectual disability and/or developmental delay. *Genome Med* 2017;**9**. doi:10.1186/s13073-017-0433-1
- 7 Sirmaci A, Spiliopoulos M, Brancati F, *et al.* Mutations in ANKRD11 cause KBG syndrome, characterized by intellectual disability, skeletal malformations, and macrodontia. *Am J Hum Genet* 2011;**89**:289–94. doi:10.1016/j.ajhg.2011.06.007
- 8 Kleyner R, Malcolmson J, Tegay D, *et al.* KBG syndrome involving a single-nucleotide duplication in ANKRD11. *Cold Spring Harb Mol Case Stud* 2016;**2**:a001131. doi:10.1101/mcs.a001131
- 9 Parenti I, Gervasini C, Pozojevic J, *et al.* Broadening of cohesinopathies: Exome sequencing identifies mutations in ANKRD11 in two patients with Cornelia de Lange-overlapping phenotype. *Clin Genet* 2016;**89**:74–81. doi:10.1111/cge.12564
- 10 van der Sluijs PJ, Jansen S, Vergano SA, *et al.* The ARID1B spectrum in 143 patients: from nonsyndromic intellectual disability to Coffin–Siris syndrome. *Genet Med* 2019;**21**:1295–307. doi:10.1038/s41436-018-0330-z
- 11 Määttänen L, Hietala M, Ignatius J, *et al.* A 69-year-old woman with Coffin–Siris syndrome. *Am J Med Genet Part A* 2018;**176**:1764–7. doi:10.1002/ajmg.a.38844
- 12 Hoyer J, Ekici AB, Ende S, *et al.* Haploinsufficiency of ARID1B, a member of the SWI/SNF-A chromatin-remodeling complex, is a frequent cause of intellectual disability. *Am J Hum Genet* 2012;**90**:565–72. doi:10.1016/j.ajhg.2012.02.007
- 13 Vals MA, Öglane-Shlik E, Nöukas M, *et al.* Coffin–Siris Syndrome with obesity, macrocephaly, hepatomegaly and hyperinsulinism caused by a mutation in the ARID1B gene. *Eur J Hum Genet* 2014;**22**:1327–9. doi:10.1038/ejhg.2014.25

- 14 Shen W, Krautscheid P, Rutz AM, *et al.* De novo loss-of-function variants of ASH1L are associated with an emergent neurodevelopmental disorder. *Eur J Med Genet* 2019;**62**:55–60. doi:10.1016/j.ejmg.2018.05.003
- 15 Wang T, Guo H, Xiong B, *et al.* De novo genic mutations among a Chinese autism spectrum disorder cohort. *Nat Commun* 2016;**7**:13316. doi:10.1038/ncomms13316
- 16 Wayhelova M, Oppelt J, Smetana J, *et al.* Novel de novo frameshift variant in the ASXL3 gene in a child with microcephaly and global developmental delay. *Mol Med Rep* 2019;**20**:505–12. doi:10.3892/mmr.2019.10303
- 17 Balasubramanian M, Willoughby J, Fry AE, *et al.* Delineating the phenotypic spectrum of Bainbridge-Ropers syndrome: 12 new patients with de novo, heterozygous, loss-of-function mutations in ASXL3 and review of published literature. *J Med Genet* 2017;**54**:537–43. doi:10.1136/jmedgenet-2016-104360
- 18 Verhoeven W, Egger J, Räkens E, *et al.* Phenotypic characterization of an older adult male with late-onset epilepsy and a novel mutation in ASXL3 shows overlap with the associated Bainbridge-Ropers syndrome. *Neuropsychiatr Dis Treat* 2018;**14**:867–70. doi:10.2147/NDT.S153511
- 19 Chinen Y, Nakamura S, Ganaha A, *et al.* Mild prominence of the Sylvian fissure in a Bainbridge-Ropers syndrome patient with a novel frameshift variant in ASXL3. *Clin Case Reports* 2018;**6**:330–6. doi:10.1002/ccr3.1361
- 20 Koboldt DC, Mosher TM, Kelly BJ, *et al.* A de novo nonsense mutation in ASXL3 shared by siblings with Bainbridge-Ropers syndrome. *Cold Spring Harb Mol Case Stud* 2018;**4**. doi:10.1101/mcs.a002410
- 21 Giri D, Rigden D, Didi M, *et al.* Novel compound heterozygous ASXL3 mutation causing Bainbridge-ropers like syndrome and primary IGF1 deficiency. *Int J Pediatr Endocrinol* 2017;**2017**. doi:10.1186/s13633-017-0047-9
- 22 Dinwiddie DL, Soden SE, Saunders CJ, *et al.* De novo frameshift mutation in ASXL3 in a patient with global developmental delay, microcephaly, and craniofacial anomalies. *BMC Med Genomics* 2013;**6**:32. doi:10.1186/1755-8794-6-32
- 23 Dias C, Estruch SB, Graham SA, *et al.* BCL11A Haploinsufficiency Causes an Intellectual Disability Syndrome and Dysregulates Transcription. *Am J Hum Genet* 2016;**99**:253–74. doi:10.1016/j.ajhg.2016.05.030
- 24 Cai T, Chen X, Li J, *et al.* Identification of novel mutations in the HbF repressor gene BCL11A in patients with autism and intelligence disabilities. *Am J Hematol* 2017;**92**:E653–6. doi:10.1002/ajh.24902
- 25 Bernardo P, Galletta D, Iasevoli F, *et al.* CHD2 mutations: Only epilepsy? Description of cognitive and behavioral profile in a case with a new mutation. *Seizure* 2017;**51**:186–9. doi:10.1016/j.seizure.2017.09.001
- 26 Yamamoto T, Imaizumi T, Yamamoto-Shimojima K, *et al.* Genomic backgrounds

- of Japanese patients with undiagnosed neurodevelopmental disorders. *Brain Dev* 2019;**41**:776–82. doi:10.1016/j.braindev.2019.05.007
- 27 Douzgou S, Liang HW, Metcalfe K, *et al.* The clinical presentation caused by truncating CHD8 variants. *Clin Genet* 2019;**96**:72–84. doi:10.1111/cge.13554
- 28 Wang J, Liu J, Gao Y, *et al.* Autism spectrum disorder early in development associated with CHD8 mutations among two Chinese children. *BMC Pediatr* 2018;**18**:338. doi:10.1186/s12887-018-1307-4
- 29 Stoleran ES, Smith B, Chaubey A, *et al.* CHD8 intragenic deletion associated with autism spectrum disorder. *Eur J Med Genet* 2016;**59**:189–94. doi:10.1016/j.ejmg.2016.02.010
- 30 Bernier R, Golzio C, Xiong B, *et al.* Disruptive CHD8 Mutations Define a Subtype of Autism Early in Development. *Cell* 2014;**158**:263–76. doi:10.1016/j.cell.2014.06.017
- 31 Merner N, Forgeot d’Arc B, Bell SC, *et al.* A de novo frameshift mutation in chromodomain helicase DNA-binding domain 8 (CHD8): A case report and literature review. *Am J Med Genet A* 2016;**170A**:1225–35. doi:10.1002/ajmg.a.37566
- 32 Codina-Solà M, Rodríguez-Santiago B, Homs A, *et al.* Integrated analysis of whole-exome sequencing and transcriptome profiling in males with autism spectrum disorders. *Mol Autism* 2015;**6**:21. doi:10.1186/s13229-015-0017-0
- 33 Chanes NM, Wong J, Lacassie Y. Further delineation of DDX3X syndrome. *Clin Dysmorphol* 2019;**28**:151–3. doi:10.1097/MCD.0000000000000263
- 34 Wang X, Posey JE, Rosenfeld JA, *et al.* Phenotypic expansion in DDX3X - a common cause of intellectual disability in females. *Ann Clin Transl Neurol* 2018;**5**:1277–85. doi:10.1002/acn3.622
- 35 Blackburn ATM, Bekheirnia N, Uma VC, *et al.* DYRK1A-related intellectual disability: a syndrome associated with congenital anomalies of the kidney and urinary tract. *Genet Med* 2019;**21**:2755–64. doi:10.1038/s41436-019-0576-0
- 36 Evers JMG, Laskowski RA, Bertolli M, *et al.* Structural analysis of pathogenic mutations in the DYRK1A gene in patients with developmental disorders. *Hum Mol Genet* 2017;**26**:519–26. doi:10.1093/hmg/ddw409
- 37 Chan AJS, Cytrynbaum C, Hoang N, *et al.* Expanding the neurodevelopmental phenotypes of individuals with de novo KMT2A variants. *npj Genomic Med* 2019;**4**:9. doi:10.1038/s41525-019-0083-x
- 38 Feldman HR, Dlouhy SR, Lah MD, *et al.* The progression of Wiedemann-Steiner syndrome in adulthood and two novel variants in the KMT2A gene. *Am J Med Genet A* 2019;**179**:300–5. doi:10.1002/ajmg.a.60698
- 39 Baer S, Afenjar A, Smol T, *et al.* Wiedemann-Steiner syndrome as a major cause of syndromic intellectual disability: A study of 33 French cases. *Clin Genet*

- 2018;**94**:141–52. doi:10.1111/cge.13254
- 40 Sun Y, Hu G, Liu H, *et al.* Further delineation of the phenotype of truncating KMT2A mutations: The extended Wiedemann-Steiner syndrome. *Am J Med Genet A* 2017;**173**:510–4. doi:10.1002/ajmg.a.38025
- 41 Alexandrou A, Papaevripidou I, Alexandrou IM, *et al.* De novo mosaic MECP2 mutation in a female with Rett syndrome. *Clin Case Reports* 2019;**7**:366–70. doi:10.1002/ccr3.1985
- 42 Castronovo P, Baccarin M, Ricciardello A, *et al.* Phenotypic spectrum of NRXN1 mono- and bi-allelic deficiency: A systematic review. *Clin Genet* 2020;**97**:125–37. doi:10.1111/cge.13537
- 43 Brignell A, St John M, Boys A, *et al.* Characterization of speech and language phenotype in children with NRXN1 deletions. *Am J Med Genet B Neuropsychiatr Genet* 2018;**177**:700–8. doi:10.1002/ajmg.b.32664
- 44 Al Shehhi M, Forman EB, Fitzgerald JE, *et al.* NRXN1 deletion syndrome; phenotypic and penetrance data from 34 families. *Eur J Med Genet* 2019;**62**:204–9. doi:10.1016/j.ejmg.2018.07.015
- 45 Sajid U, Argiropoulos B, Wei X-C, *et al.* Two De Novo Mutations in an Autistic Child Who Had Previously Undergone Transplantation for Dilated Cardiomyopathy: The Importance of Keeping an Open Mind. *Can J Cardiol* 2017;**33**:292.e5–292.e7. doi:10.1016/j.cjca.2016.08.019
- 46 Ferretti A, Barresi S, Trivisano M, *et al.* POGZ-related epilepsy: Case report and review of the literature. *Am J Med Genet A* 2019;**179**:1631–6. doi:10.1002/ajmg.a.61206
- 47 Samanta D, Ramakrishnaiah R, Schaefer B. The neurological aspects related to POGZ mutation: case report and review of CNS malformations and epilepsy. *Acta Neurol. Belg.* 2019. doi:10.1007/s13760-019-01122-6
- 48 Du X, Gao X, Liu X, *et al.* Genetic Diagnostic Evaluation of Trio-Based Whole Exome Sequencing Among Children With Diagnosed or Suspected Autism Spectrum Disorder. *Front Genet* 2018;**9**. doi:10.3389/fgene.2018.00594
- 49 Dentici ML, Niceta M, Pantaleoni F, *et al.* Expanding the phenotypic spectrum of truncating POGZ mutations: Association with CNS malformations, skeletal abnormalities, and distinctive facial dysmorphism. *Am J Med Genet A* 2017;**173**:1965–9. doi:10.1002/ajmg.a.38255
- 50 Stessman HAF, Willemsen MH, Fenckova M, *et al.* Disruption of POGZ Is Associated with Intellectual Disability and Autism Spectrum Disorders. *Am J Hum Genet* 2016;**98**:541–52. doi:10.1016/j.ajhg.2016.02.004
- 51 White J, Beck CR, Harel T, *et al.* POGZ truncating alleles cause syndromic intellectual disability. *Genome Med* 2016;**8**. doi:10.1186/s13073-015-0253-0
- 52 Ye Y, Cho MT, Retterer K, *et al.* De novo POGZ mutations are associated with

- neurodevelopmental disorders and microcephaly . *Mol Case Stud* 2015;**1**:a000455. doi:10.1101/mcs.a000455
- 53 Fukai R, Hiraki Y, Yofune H, *et al.* A case of autism spectrum disorder arising from a de novo missense mutation in POGZ. *J Hum Genet* 2015;**60**:277–9. doi:10.1038/jhg.2015.13
- 54 Nickel K, Tebartz van Elst L, Domschke K, *et al.* Heterozygous deletion of SCN2A and SCN3A in a patient with autism spectrum disorder and Tourette syndrome: a case report. *BMC Psychiatry* 2018;**18**:248. doi:10.1186/s12888-018-1822-8
- 55 Leach EL, van Karnebeek CDM, Townsend KN, *et al.* Episodic ataxia associated with a de novo SCN2A mutation. *Eur J Paediatr Neurol* 2016;**20**:772–6. doi:10.1016/j.ejpn.2016.05.020
- 56 Wong VCN, Fung CW, Kwong AKY. SCN2A mutation in a Chinese boy with infantile spasm - response to Modified Atkins Diet. *Brain Dev* 2015;**37**:729–32. doi:10.1016/j.braindev.2014.10.008
- 57 Tavassoli T, Kolevzon A, Wang AT, *et al.* De novo SCN2A splice site mutation in a boy with Autism spectrum disorder. *BMC Med Genet* 2014;**15**:35. doi:10.1186/1471-2350-15-35
- 58 Baasch AL, Hüning I, Gilissen C, *et al.* Exome sequencing identifies a de novo SCN2A mutation in a patient with intractable seizures, severe intellectual disability, optic atrophy, muscular hypotonia, and brain abnormalities. *Epilepsia* 2014;**55**. doi:10.1111/epi.12554
- 59 Powis Z, Farwell Hagman KD, Mroske C, *et al.* Expansion and further delineation of the SETD5 phenotype leading to global developmental delay, variable dysmorphic features, and reduced penetrance. *Clin Genet* 2018;**93**:752–61. doi:10.1111/cge.13132
- 60 Grozeva D, Carss K, Spasic-Boskovic O, *et al.* De novo loss-of-function mutations in SETD5, encoding a methyltransferase in a 3p25 microdeletion syndrome critical region, cause intellectual disability. *Am J Hum Genet* 2014;**94**:618–24. doi:10.1016/j.ajhg.2014.03.006
- 61 Li Y, Jia X, Wu H, *et al.* Genotype and phenotype correlations for SHANK3 de novo mutations in neurodevelopmental disorders. *Am J Med Genet Part A* 2018;**176**:2668–76. doi:10.1002/ajmg.a.40666
- 62 Kanani F, Study D, Balasubramanian M. SHANK3 variant as a cause of nonsyndromal autism in an 11-year-old boy and a review of published literature. *Clin Dysmorphol* 2018;**27**:113–5. doi:10.1097/MCD.000000000000232
- 63 Zhu W, Li J, Chen S, *et al.* Two de novo novel mutations in one SHANK3 allele in a patient with autism and moderate intellectual disability. *Am J Med Genet A* 2018;**176**:973–9. doi:10.1002/ajmg.a.38622
- 64 Terrone G, Vitiello G, Genesisio R, *et al.* A novel SHANK3 interstitial microdeletion in a family with intellectual disability and brain MRI abnormalities resembling

- Unidentified Bright Objects. *Eur J Paediatr Neurol* 2017;**21**:902–6. doi:10.1016/j.ejpn.2017.07.006
- 65 Cook EH, Masaki JT, Guter SJ, *et al.* Lovastatin Treatment of a Patient with a De Novo SYNGAP1 Protein Truncating Variant. *J Child Adolesc Psychopharmacol* 2019;**29**:321–2. doi:10.1089/cap.2018.0159
- 66 von Stülpnagel C, Hartlieb T, Borggräfe I, *et al.* Chewing induced reflex seizures ('eating epilepsy') and eye closure sensitivity as a common feature in pediatric patients with SYNGAP1 mutations: Review of literature and report of 8 cases. *Seizure* 2019;**65**:131–7. doi:10.1016/j.seizure.2018.12.020
- 67 Pei Y, Li W, Du L, *et al.* Novel Mutation of SYNGAP1 Associated with Autosomal Dominant Mental Retardation 5 in a Chinese Patient. *Fetal Pediatr Pathol* 2018;**37**:400–3. doi:10.1080/15513815.2018.1497113
- 68 Kimura Y, Akahira-Azuma M, Harada N, *et al.* Novel SYNGAP1 variant in a patient with intellectual disability and distinctive dysmorphisms. *Congenit Anom (Kyoto)* 2018;**58**:188–90. doi:10.1111/cga.12273
- 69 Prchalova D, Havlovicova M, Sterbova K, *et al.* Analysis of 31-year-old patient with SYNGAP1 gene defect points to importance of variants in broader splice regions and reveals developmental trajectory of SYNGAP1-associated phenotype: case report. *BMC Med Genet* 2017;**18**:62. doi:10.1186/s12881-017-0425-4
- 70 Berryer MH, Hamdan FF, Klitten LL, *et al.* Mutations in SYNGAP1 cause intellectual disability, autism, and a specific form of epilepsy by inducing haploinsufficiency. *Hum Mutat* 2013;**34**:385–94. doi:10.1002/humu.22248
- 71 Vetrini F, McKee S, Rosenfeld JA, *et al.* De novo and inherited TCF20 pathogenic variants are associated with intellectual disability, dysmorphic features, hypotonia, and neurological impairments with similarities to Smith–Magenis syndrome. *Genome Med* 2019;**11**:12. doi:10.1186/s13073-019-0623-0
- 72 Torti E, Keren B, Palmer EE, *et al.* Variants in TCF20 in neurodevelopmental disability: description of 27 new patients and review of literature. *Genet Med* 2019;**21**:2036–42. doi:10.1038/s41436-019-0454-9
- 73 Schäffgen J, Cremer K, Becker J, *et al.* De novo nonsense and frameshift variants of TCF20 in individuals with intellectual disability and postnatal overgrowth. *Eur J Hum Genet* 2016;**24**:1739–45. doi:10.1038/ejhg.2016.90