

CHARGE syndrome: the phenotypic spectrum of mutations in the *CHD7* gene

Jongmans MCJ¹, Admiraal RJ², van der Donk KP¹, Vissers LELM¹, Baas AF¹, Kapusta L³, van Hagen JM⁴, Donnai D⁵, de Ravel TJ⁶, Veltman JA¹, Geurts van Kessel A¹, De Vries BBA¹, Brunner HG¹, Hoefsloot LH¹, van Ravenswaaij CMA¹

- 1) Departments of Human Genetics, 2) Otorhinolaryngology and 3) Children's Heart Centre, Radboud University Nijmegen Medical Centre, Nijmegen, the Netherlands.
- 4) Department of Clinical Genetics and Human Genetics, VU University Medical Centre, Amsterdam, the Netherlands
- 5) Academic Unit of Medical Genetics and Regional Genetics Service, St. Mary's Hospital, Manchester, United Kingdom.
- 6) Centre for Human Genetics, UZ Gasthuisberg, University of Leuven, Leuven, Belgium.

Key words: CHARGE syndrome, *CHD7*, clinical spectrum

Correspondence:

C. van Ravenswaaij
Dept. of Human Genetics,
Radboud University Nijmegen Medical Centre
P.O. Box 9101
6500 HB Nijmegen The Netherlands
Phone 31-24-3613946 Fax 31-24-3668753
e-mail c.vanravenswaaij@antrg.umcn.nl

ABSTRACT

Background: CHARGE syndrome is a non-random clustering of congenital anomalies. The *CHD7* gene on chromosome 8q12.1 was recently discovered as a major gene involved in the aetiology of this syndrome.

Methods: The coding regions of *CHD7* were screened for mutations in 107 index patients with clinical features suggestive of CHARGE syndrome. Clinical data of the mutation positive patients were sampled to study the phenotypic spectrum of mutations in the *CHD7* gene.

Results: Mutations were identified in 69 patients. Here we describe the clinical features of 47 of these patients, including two sib pairs. Most mutations were unique and were scattered throughout the gene. All patients but one fulfilled the current diagnostic criteria for CHARGE syndrome. No genotype-phenotype correlations were apparent in this cohort, which is best demonstrated by the differences in clinical presentation in sib pairs with identical mutations. Somatic mosaicism was detected in the unaffected mother of a sib pair, supporting the existence of germline mosaicism.

Conclusions: *CHD7* mutations account for the majority of the cases with CHARGE syndrome, with a broad clinical variability and without an obvious genotype-phenotype correlation. In one case evidence for germline mosaicism was provided.

INTRODUCTION

CHARGE syndrome (OMIM #214800) is a pleiotropic disorder comprising of coloboma, hear defects, choanal atresia, retarded growth and development, genital hypoplasia, ear anomalies and deafness. A consistent feature in CHARGE syndrome is semicircular canal hypoplasia resulting in vestibular areflexia.¹⁻³ Other commonly associated congenital anomalies are facial nerve palsy, cleft lip/palate and tracheo-esophageal fistula. Specific behavioural problems, including autistic-like, have been described.^{4,5} The combination of abnormalities initially known as CHARGE association was first reported independently by Hall and by Hittner and colleagues in 1979,^{6,7} after which Pagon and colleagues proposed the acronym CHARGE in 1981.⁸ CHARGE syndrome is an autosomal dominant syndrome with an estimated prevalence at birth between 1 per 10,000 to 1 per 15,000.⁹ Recent epidemiological data revealed an occurrence of CHARGE syndrome in 1 in 8,500 live births in the Atlantic Provinces of Canada.¹⁰

CHARGE syndrome is a phenotypically heterogeneous syndrome and its clinical diagnosis is made using criteria that have been refined several times. Blake et al. suggested diagnostic criteria in 1998.⁹ A refinement of these criteria for different age groups was proposed to capture the continuum of the presentation of CHARGE syndrome.¹⁰ Simultaneously, Verloes suggested an update of diagnostic criteria, emphasizing the most specific embryological defects while avoiding non-specific or secondary anomalies.¹¹ He also suggested the exclusion of sex-dependent criteria. Both sets of diagnostic criteria are depicted in Table I. CHARGE syndrome was only recently reconsidered to be a syndrome, instead of an association since our group discovered *CHD7* on chromosome 8 (8q12.1) as a major gene involved in this syndrome.¹² *CHD7* encodes a protein of the chromodomain (chromatin organization modifier) family. Members of this family share a unique combination of functional domains consisting of two N-terminal chromodomains, followed by a SWI2/SNF2-like ATPase/helicase domain and a DNA binding domain.¹³⁻¹⁴ It is assumed that CHD protein complexes affect chromatin structure and gene expression and, thereby, play an important role in regulating embryonic development.

We report a study of the phenotypic spectrum in 47 patients with a *CHD7* mutation, with special emphasis on differences in presentation in sib pairs that share identical mutations.

Table I. Updated diagnostic Criteria for CHARGE syndrome

Verloes 2005 Typical CHARGE = 3 major criteria or 2 major and 2 minor criteria

Major Criteria

- 1) Coloboma (iris or choroid, with or without microphthalmia)
- 2) Atresia of Choanae
- 3) Hypoplastic semicircular Canals

Minor criteria

- 1) Rhombencephalic dysfunction (brainstem dysfunctions, cranial nerve VII to XII palsies and neurosensory deafness)
- 2) Malformation of mediastinal organs (heart, esophagus)
- 3) Hypothalamo-hypophyseal dysfunction (including GH and gonadotrophin deficiencies)
- 4) Abnormal middle or external ear
- 5) Mental retardation

Blake et al. 1998 All four major criteria, or three major and three minor criteria

Major criteria

- 1) Coloboma- iris, retina, choroid, optic disc or microphthalmia
- 2) Atresia of Choanae
- 3) Cranial nerve dysfunction I: Anosmia, VII:Facial palsy, VIII Sensorineural deafness and vestibular problems, IX and/or X: Swallowing problems
- 4) Characteristic external ears (absent or hypoplastic lobes, asymmetry, decreased cartilaginous folds, and triangular concha) and inner ear anomalies (temporal bone findings with cochlear hypoplasia and or absent/hypoplastic semicircular canals)

Minor criteria

- 1) Characteristic facial features - broad, sloping forehead, laterally protruding ears, small mouth, and high nasal bridge
- 2) Congenital cardiovascular malformations of all types
- 3) Tracheo-oesophageal fistula
- 4) Growth deficiency
- 5) Genital hypoplasia - micropenis and/or cryptorchidism or hypoplastic labia. Delayed, incomplete pubertal development
- 6) Orofacial cleft
- 7) Developmental delay: delayed motor milestones, hypotonia, MR

METHODS

Patients

The coding regions of the *CHD7* gene were screened for mutations in 107 index patients with clinical features suggestive of CHARGE syndrome. In 69 of these patients a mutation was identified (65 %) and for 47 patients (22 males, 25 females, two sib pairs) sufficient clinical data was available to include them for further studies. The cohort includes fifteen patients reported in our previous study.¹² Parental DNA samples of 22 patients, including one sib pair, were tested for *de novo* occurrence.

Clinical information of the patients was obtained through investigation in our own department or through a written questionnaire when DNA of the patient was referred to the DNA-diagnostics section of our department. Additional information was obtained from clinicians when necessary. The diagnostic criteria by Blake and Verloes (table I) were applied to all cases for which sufficient clinical information was available.^{9, 11}

All patients or their legal representatives gave informed consent for the DNA studies and the collection of clinical data.

Mutation screening

DNA was isolated according to standard procedures. The 37 coding exons of the *CHD7* gene (exon 2-38, accession number NM_017780) and their flanking intron sequences were amplified by polymerase chain reaction (PCR). Subsequently, sequence analysis was performed using a 3730 automated sequencer (Applied Biosystems).

The primer sets used previously were optimized by using shorter PCR products to exclude allele dropout.¹² Primer information and PCR conditions are given in supplemental tables I and II, available from the *JMG* website.

Whole gene deletions were excluded by Multiplex Ligation-dependent Probe Amplification (MLPA). Specific probe sets were designed for exons 2-11 and exons 33-38. MLPA analysis was performed according to the instructions of the manufacturer (MRC Holland b.v.; www.mlpa.com). Probe information is given in table III of the supplemental data (*JMG* website).

RESULTS

CHD7 mutation analysis

Mutation analysis in our series of 107 index patients revealed 69 mutations in the *CHD7* gene (Fig. 1 and Table II). Two mutations were recurrent, all others were unique. We detected 31 nonsense, 17 frame shift, 13 splice site and 8 missense mutations that were scattered throughout the gene. In the affected sibs identical mutations were identified. Among the *CHD7* positive patients is one girl (no. 22) with a previously identified chromosome 22q11 deletion. Fifteen patients were reported in a previous study.¹² In six of these patients, however, the mutation was not detected initially. After a more thorough investigation with improved primer sets *CHD7* mutations were detected. For 21 index patients the parents were studied. In 20 cases the mutation was proven to be *de novo*. In the sib pair consisting of two boys, mosaicism for the *CHD7* mutation was identified in the mother. In the remaining 38 mutation-negative patients whole gene deletions were excluded by MLPA analysis.

Table II. Overview of *CHD7* mutations

Mutation in <i>CHD7</i>	exon	Theoretical effect on RNA(r.) or protein (p.) *	segregation	Pt. in Table III
c.77_78delAA	2	p.Glu26fs	nd	1
c.469C>T	2	p.Arg157X	nd	
c.921_922delAG (e2)	2	p.Gly308fs	nd	
c.1044delC	2	p.Asn349fs	nd	2
c.1078G>T	2	p.Gly360X	one parent excl.	3
c.1388delG	2	p.Gly463fs	de novo	4
c.1465C>T	2	p.Gln489X	nd	5
c.1495C>T	2	p.Gln499X	nd	
c.1714C>T	3	p.Gln572X	de novo	6
c.1973_1974insT	3	p.Glu658fs	nd	7
c.2095A>G	3	r.spl? p.S699G	nd	
c.2194C>G	4	p.Pro732Ala	nd	
c.2238+1G>A	IVS4	r.spl?	de novo	8
c.2442+5G>A	IVS6	r.spl?	nd	9
c.2504_2508delATCTT	8	p.Tyr835fs	nd	
c.2505T>A	8	p.Tyr835X	de novo	10
c.2520G>A	8	p.Trp840X	nd	
c.2572C>T	8	p.Arg858X	de novo	11
c.2958-2A>T	IVS11	r.spl?	de novo	12
c.2959C>T	12	p.Arg987X	nd	13
c.3053_3054insA	12	p.Phe1019fs	nd	14
c.3082A>G	12	p.Ile1028Val	de novo	45
c.3106C>T	12	p.Arg1036X	nd	
c.3302G>A	13	p.Cys1101Tyr	nd	
c.3654C>G	15	p.Tyr1218X	nd	
c.3655C>T	15	p.Arg1219X	nd	15
c.3770T>G	15	p.Leu1257Arg	de novo	46
c.3779-2A>G	IVS15	r.spl?	nd	16
c.4015C>T	17	p.Arg1339X	nd	
c.4157C>G	17	p.Ser1386X	one parent excl.	17
c.4226_4227delTG	18	p.Val1409fs	nd	18
c.4507G>T	19	p.Glu1503X	nd	19

c.4644+1G>A	IVS20	r.spl?		nd	
c.4787A>G	21	p.Asp1596Gly		nd	
c.5050-41_5050-3del39	IVS23	r.spl?		nd	20
c.5402A>C	25	p.His1801Pro		de novo	47
c.5405-17G>A	IVS25	r.spl?		de novo	21
c.5405-7G>A	IVS25	r.spl?		nd	
c.5418C>G	26	p.Asn1807X	22q11del	de novo	22
c.5436C>A	26	p.Asp1812Glu		nd	
c.5534G>A	IVS26	r.spl?		de novo	23
c.5668A>T	29	p.Lys1890X		de novo	24
c.5680_5681delAG	29	p.Ser1894fs		de novo	25
c.5752_5753dupA	29	p.Thr1918fs	sib pair 1	nd	26
c.5752_5753dupA	29	p.Thr1918fs	sib pair 1	nd	27
c.5833C>T	29	p.Arg1945X		de novo	28
c.5893+1G>A	IVS29	r.spl?		nd	
c.5982G>A	30	p.Trp1994X	#	nd	
c.5982G>A	30	p.Trp1994X	#	sib pair 2	mat mosaicism 29
c.5982G>A	30	p.Trp1994X	#	sib pair 2	mat mosaicism 30
c.6051T>A	30	p.Cys2017X		nd	31
c.6070C>T	30	p.Arg2024X		one parent excl.	32
c.6079C>T	30	p.Arg2027X		de novo	33
c.6148C>T	31	p.Arg2050X	\$	nd	35
c.6148C>T	31	p.Arg2050X	\$	nd	34
c.6155_6157CTC>AGA	31	p.Ser2052X		nd	
c.6157C>T	31	p.Arg2053X		nd	
c.6304delG	31	p.Val2102fs		de novo	36
c.6775+2_6775+3insGT	IVS31	r.spl?		nd	37
c.6955C>T	33	p.Arg2319Cys		nd	
c.7079delA	33	p.Lys2360fs		nd	
c.7165-4A>G	IVS33	r.spl? p.Lys2388_Glu2389insX		nd	
c.7180delC	34	p.Lys2394fs		nd	38
c.7219delA	34	p.Ile2407fs		nd	
c.7252C>T	34	p.Arg2418X		nd	39
c.7400delT	34	p.Leu2467fs		de novo	40
c.7824T>A	35	p.Tyr2608X		de novo	41
c.7879C>T	36	p.Arg2627X		nd	42
c.7884_7885delTA	36	p.His2628fs		de novo	43
c.8016G>A	37	p.Trp2672X		nd	
c.8744_8745dupG	38	p.Leu2916fs		nd	44

* nomenclature according to <http://www.genomic.unimelb.edu.au/mdi/mutnomen/> "

nd = not done

#\$ = recurrent mutation

Clinical features

Information obtained through our own investigation and/or through written questionnaires, supplemented with additional information from clinicians, resulted in clinical features of the 47 selected patients as outlined in Table III. Details of these features are provided below. All 47 cases were included in the evaluation unless stated otherwise. The diagnostic criteria by Blake and Verloes (table I) could be applied to 38 cases.^{9 11} Only one patient did not fulfil both sets of diagnostic criteria.

Sufficient clinical information could also be obtained for 23 out of the 38 *CHD7* negative patients. Of these patients only two fulfilled the clinical diagnostic criteria by Blake and Verloes.^{9 11}

In this section first a summary of all clinical data of the 47 *CHD7* positive patients is given. Subsequently a detailed case report is provided of a girl who did not fulfil the diagnostic criteria by Blake and Verloes (table I)^{9 11} and the intra familial variability in sib pairs is delineated.

Neonatal period

The median gestational age of the patients was 38.2 weeks (n=45, range 30-42 weeks). Only one patient was reported to be small for gestational age, while feeding difficulties were reported in 33 (70%) patients. Four patients required a gastrostomy due to severe feeding problems.

Four patients died during the neonatal period, three during the first half year of life and one patient at the age of fourteen years. At the time of investigation four patients were below the age of one year.

Coloboma of the eye

In 33 patients (70%) a coloboma of one (n=4) or both (n=29) eyes was present. In only nine patients the iris was involved, thus in the majority the coloboma was only visible by fundoscopy. In none of the patients the coloboma was restricted to the iris only.

Microphthalmia was present in ten patients (21%).

Congenital heart defects

Thirty-one (66%) patients had a congenital heart defect. Fourteen (30%) patients had major heart defects: six tetralogy of Fallot, two double-outlet right ventricle (one combined with hypoplastic left heart and AVSD), three isolated hypoplastic left heart syndrome, one hypoplastic right heart syndrome, one agenesis of the pulmonary valve combined with hypoplastic left heart and one Shone's complex. A right descending aorta was present in three patients and one patient had a vascular ring. The other patients had solitary patent ductus arteriosus beyond infancy (3), patent ductus arteriosus combined with atrium septum defect and/or ventricular septum defect (6) or a solitary septal defect (4).

Retardation of growth and development

A height below the third percentile was reported in 21 out of 32 patients (63%).

Speech development varied from mild speech delay to a severe retardation without speech. Learning disabilities were reported in 24 (75%) out of 32 patients who were above the age of twelve months at last examination. Eight patients (25%) had no cognitive impairment.

Endocrine and urogenital abnormalities

At the moment of *CHD7* testing fifteen patients (8 girls, seven boys) were above fifteen years of age. Gonadotrophin deficiency was present in seven (88%) of these girls, and six (86%) of these boys. Two girls had their menarche at age 14. A hypoplastic uterus was found by ultrasound investigation in three girls. Of all 22 mutation-positive boys four (18%) had cryptorchidism, six (27%) micropenis and seven (32%) had both cryptorchidism and micropenis.

Three patients had a horseshoe kidney and in two patients agenesis of the left kidney was demonstrated. A vesicoureteral reflux was reported in three patients and one patient had renal cysts.

Table III. Anomalies in 47 CHD7 positive patients

Individual	Sex	Age at clinical evaluation	Gestation (weeks)	Birth weight (grams)	Coloboma		Microphthalmia	Heart defect	Atresia of choanae	Height	Micropenis / Cryptorchidism	Ext. ear anomaly	Hearing loss	Vestibular dysfunction	Facial nerve palsy	Oesophageal fistula	Cleft lip	Cleft palate
					Left	Right												
1	F	17	38	2450	IRCO	IRCO	L/R	-	-	<P3	+	+	?	-	-	-	-	
2	F	19	41	3520	RC	RC	R	+	-	?	+	+	?	-	-	-	+	
3	F	17	38	2435	-	-	-	+	+	<P3	+	-	+++	-	-	-	-	
4	F	6	32	1805	-	-	-	+	-	<P3	+	+	+++	-	+	+	+	
5	F	6	36	?	IRCO	IRCO	-	+	-	?	+	?	?	-	-	+	+	
6	M	10	38	3025	-	-	-	-	-	P3	+	+	+	+++	+	-	-	+
7	F	11	40	3350	CRO	CRO	L/R	-	-	?	+	+	+++	-	-	-	+	
8	M	7	39	2530	CRO	O	L	-	-	<P3	+	+	+	+++	-	-	-	-
9	M	4	39	3200	-	-	-	-	-	>P10	+	+	+	?	-	-	+	+
10	M	<1	31	1650	-	-	-	-	+	?	+	+	+	+++	-	-	-	-
11	F	<1	40	4082	R	R	-	+	-	?	-	+	+	?	-	-	+	+
12	M	day 6 ^A	38	3060	IRO	RO	-	+	-	?	-	+	?	?	-	-	+	+
13	M	32	37	2900	-	-	-	-	+	<P3	+	+	+	?	-	-	-	-
14	F	35	?	3250	C	-	L	+	+	<P3	+	+	+++	-	-	-	-	-
15	F	15	39	2910	-	-	-	+	+	<P3	+	+	++	-	-	-	-	-
16	M	19	42	3700	IRC	IRC	-	-	-	P15	+	+	+	++	-	-	-	-
17	M	19	40	2870	-	-	-	+	+	<P3	+	+	+	+	-	-	-	-
18	M	day 12 ^A	35	2250	IO	O	-	+	+	?	+	+	+	+++	-	-	-	-
19	F	5	40	3840	RO	RO	-	+	-	<P3	+	-	?	-	-	+	+	
20	M	month 5 ^A	38	3408	-	-	-	+	+	?	+	?	?	-	-	-	-	-
21	M	40	40	3250	RO	RO	R	+	+	P10	+	+	+	?	+	+	-	-
22	F	22	36	2000	-	-	-	+	+	<P3	+	+	+	+++	-	+	-	-
23	M	19	42	2800	IR	R	-	+	-	<P3	+	+	+	+++	-	-	+	+
24	F	month 6 ^A	42	3450	IR	IR	-	+	-	?	-	+	+	?	+	-	-	-
25	M	1	37	2940	R	R	-	+	-	P50	+	+	+	+++	+	+	-	-
26	F	11	35	1910	RC	IRC	-	+	+	<P3	+	+	+	+	-	-	-	-
27	F	day 2 ^A	35	1500	-	-	-	+	+	?	+	?	?	-	+	-	-	-
28	F	5	40	3700	-	-	-	+	-	P3	+	+	+++	-	-	-	-	-
29	M	7	40	2867	-	-	-	+	-	<P3	+	+	-	?	+	-	+	+
30	M	3	38	3440	RC	-	-	-	-	P3	+	+	+	?	-	+	-	-
31	M	year 14 ^A	?	3500	RC	RC	-	-	-	<P3	-	+	+	+++	-	+	+	+
32	F	12	41	2800	IR	R	L	+	-	<P10	+	+	+	+++	+	-	-	-
33	M	week 5 ^A	30	1470	RO	RO	U	-	+	?	-	+	+	?	-	+	-	-
34	M	6	37	2700	R	R	-	+	-	?	-	+	+	?	-	-	-	+
35	M	6	35	2835	O	R	R	+	-	<P3	+	+	+	+++	+	-	-	+
36	M	20	40	2820	R	R	-	-	+	<P3	+	+	+	?	-	-	+	-
37	M	<1	39	3114	R	R	-	+	-	?	+	+	?	?	-	-	-	-
38	F	day 21 ^A	35	2393	CRO	CRO	-	+	+	?	+	?	?	-	-	-	-	-
39	F	3	40	3340	CR	CR	-	+	-	<P3	+	+	+	+++	-	-	-	-
40	M	20	42	3490	CR	CR	-	+	+	<P3	+	+	+	+	-	-	+	+
41	F	10	37	2650	-	-	-	+	+	<P3	+	+	+	+++	-	-	-	-
42	F	<1	41	2830	CO	O	-	+	-	?	+	+	?	?	+	-	-	-
43	F	26	40	2450	R	-	-	+	-	<P3	+	+	+++	+	-	-	-	-
44	F	20	36	2450	CRO	CRO	-	-	+	?	+	+	+++	+	-	+	+	+
Missense mutations:																		
45	F	15	38	2980	O	O	L/R	-	-	<P3	+	+	+++	-	-	-	-	-
46	F	15	41	2500	C	C	-	-	-	P10	+	+	+++	-	-	-	-	-
47	F	16	40	2100	C	-	-	-	+	P3	+	-	+	-	-	-	-	-

^A= deceased

Coloboma: I = iris, R = retina, C = choroidea, O = optic disc

Microphthalmia: R = right, L = Left, U = unilateral, side not known

Vestibular dysfunction: +++ = semicircular canal agenesis on CT scan of inner ear, ++ = vestibular areflexia, + = history of unsteadiness

Individual	Sex	Age at clinical evaluation	Mental Retardation	Neurological abnormalities	Skeletal Abnormalities	Urogenital findings	Gonadotrophin deficiency	Diagnostic criteria (table 1)	
								Blake (1998) ⁹	Verloes (2005) ¹¹
1	F	17	+++				+	+	
2	F	19	?				+		
3	F	17	+				+	+	
4	F	6	+					+	+
5	F	6	+++						
6	M	10	-					+	+
7	F	11	++		S			+	+
8	M	7	+++			HK		+	+
9	M	4	?						
10	M	<1	?					+	+
11	F	<1	?			C		+	
12	M	day 6 ^A	?	AC + CH	S + HV		+		
13	M	32	+++		HV		+	+	
14	F	35	++	C	S		+	+	+
15	F	15	+					+	+
16	M	19	-		S		+	+	+
17	M	19	++	ACC	S	R	+	+	+
18	M	day 12 ^A	?					+	+
19	F	5	+	H				+	
20	M	month 5 ^A	?	C		A			
21	M	40	+++				+	+	+
22	F	22	++	H + C		R	+	+	+
23	M	19	-					+	+
24	F	month 6 ^A	?						
25	M	1	?					+	+
26	F	11	++					+	+
27	F	day 2 ^A	?						
28	F	5	+					-	-
29	M	7	++					+	
30	M	3	+++					+	
31	M	year 14 ^A	++					+	+
32	F	12	++		K			+	+
33	M	week 5 ^A	?						+
34	M	6	?						
35	M	6	?			HK	+	+	+
36	M	20	+++	C			+	+	+
37	M	<1	?		T	R			
38	F	day 21 ^A	?			HK			+
39	F	3	-			A		+	+
40	M	20	+++		S		+	+	+
41	F	10	+++					+	+
42	F	<1	?						
43	F	26	-				+	+	+
44	F	20	-				+	+	+
Missense mutations									
45	F	15	-					+	+
46	F	15	+					+	+
47	F	16	-	C	HV			+	

^A = deceased

Mental Retardation: - = normal intelligence, + = mild MR, ++ = moderate MR, +++ = severe MR
Neurological abnormalities: AC = agenesis of corpus callosum, CH = cerebellar hypoplasia, C = convulsions, ACC = atrophy of cerebral cortex, H = hydrocephaly,
Skeletal abnormalities: S = scoliosis, HV = hypoplastic vertebrae, K = kyphosis, T = triphalangeal thumb
Urogenital anomalies: HK = horseshoe kidney, C = renal cysts, R = reflux, A = agenesis of one kidney

Ear and vestibular abnormalities

In all patients dysmorphisms of the ears were noted, ranging from “typical CHARGE ears” (small, square, low-set and protruding) to minor structural abnormalities such as absence of an earlobe. One patient had a pre-auricular pit and one patient had narrow external auditory canals. Hearing impairment was demonstrated in 37 out of 41 patients (90%). In 27 patients severe bilateral hearing impairment was observed, whereas five patients showed asymmetric hearing impairment with unilateral normal or mild hearing loss.

In all 21 patients who underwent CT-scanning of the temporal bones, agenesis of the semicircular canals was demonstrated. Vestibular areflexia was demonstrated in two more patients and four patients had a history of balance disturbances. This results in 27 patients (57%) with some evidence of vestibular anomaly. However, from the remaining patients information on this subject was not available, although motor delay (possibly due to vestibular areflexia) was present in all cases on direct questioning.

Nasopharyngeal abnormalities and clefting

Choanal atresia was present in 17 patients (36%) and was unilateral in only three of them.

Respiratory insufficiency during the neonatal period was reported in 24 patients (51%).

Twenty-two of them had either choanal atresia or a congenital heart defect or both.

Tracheomalacia was present in one patient.

Clefting was present in 17 patients (36%). Eleven patients had a cleft lip and palate, five had an isolated cleft palate and one had an isolated cleft lip.

Gastrointestinal abnormalities

Eight patients (17%) had esophageal atresia, which in three was accompanied by a tracheo-esophageal fistula. Two patients had a diaphragmatic hernia and one anal stenosis.

Neurological abnormalities

A minority of the patients (n=4, 9%) had central nervous system abnormalities, including corpus callosum agenesis combined with cerebellar hypoplasia (n=1), hydrocephaly (n=2), and atrophy of the cerebral cortex (n=1). Five patients had convulsions.

Facial nerve palsy was present in ten patients (21%) and mostly (9 out of 10) involved the right-sided facial nerve.

Skeletal abnormalities

Scoliosis was demonstrated in six patients (13%), kyphosis in one, abnormalities of the vertebral bodies in three (6%) and in one patient a triphalangeal thumb was demonstrated.

Aspecific CHARGE syndrome

Patient 28 (born at 40 weeks' gestation; birth weight 3700 g, 70th centile, Fig 2) was a 5 years old girl with developmental delay, slightly dysmorphic ears and severe hearing impairment. CT scan showed bilateral agenesis of the semicircular canals. She required a gastrostomy due to severe feeding problems and she had surgery on a congenital vascular ring. Her height was at the third centile. No choanal atresia, nor cleft palate or coloboma could be detected. In this girl, the only individual in our *CHD7* positive series who did not fulfil the current diagnostic criteria for CHARGE syndrome (table I)^{9 11} a *de novo* nonsense mutation was identified, 5833C>T (R1945X) in exon 29 of *CHD7*.

Familial cases

Two sib pairs were included from two families. In both cases, identical *CHD7* mutations were identified in the two sibs. Interestingly, in both cases the affected sib pairs showed distinct clinical features.

Sib pair 1 represented monozygotic twin sisters (born at 35 weeks' gestation, Table III: patients 26+27, Figs. 3a and 3b). Patient 27, who had a birth weight of 1500 g (5th-10th centile), died 29 hours after birth due to the combination of a hypoplastic left heart syndrome

and bilateral choanal atresia. Furthermore, she had a tracheo-esophageal fistula and typical "CHARGE ears". A hearing test was not performed. There were no colobomata of the irides. Patient 26 had a birth weight of 1910 g (25th centile). When examined at the age of 12 years, she had short stature (< 3rd centile) and was functioning four years behind her chronological age. She was born with a large patent ductus arteriosus that required surgery and she needed numerous procedures to correct bilateral choanal atresia. Her first years of life were complicated by feeding problems, for which she had a gastrostomy until the age of six years. She had severe bilateral deafness, abnormal external ears like her twin sister and bilateral chorioretinal colobomata with a right-sided iris coloboma and an unusual inferior pigment pattern in her left iris. Agenesis of the semicircular canals was not tested for by CT-scan, but her gait was unsteady.

Zygosity testing with five unlinked markers was performed and the results were consistent with the twins being monozygotic. In both sisters the spectrum of congenital anomalies was caused by an insertion 5752_5753insA in exon 29 of *CHD7*.

Sib pair 2 consisted of two brothers (Table III: patients 29+30, Figs. 3c and 3d). Patient 29 was 7 years of age (born at 40 weeks' gestation; birth weight 2867 g, 10th centile). He had surgery for cleft lip/palate and a complex heart defect (DORV, AVSD, hypoplastic left heart). He had short stature (3rd centile) and severe developmental delay. His ears showed the typical CHARGE dysmorphisms and he had bilateral hearing loss and unilateral facial nerve palsy. He had no colobomata or choanal atresia.

Patient 30, who was 4 years younger, had bilateral hearing loss and typical "CHARGE ears", a coloboma of the left retina and choroid and he underwent surgery on a tracheo-esophageal fistula. He had short stature (<3rd centile) and was severely mentally retarded. He had vocal cord palsy. This boy had no heart defect and no choanal atresia.

Both brothers have a 5982G>A (W1994X) mutation in exon 30 of *CHD7*. Sequence analysis of both parents revealed no mutation in the father and a minor aberrant peak in DNA extracted from lymphocytes of the mother. This indicated that a possible mosaicism was present in the mother. This was further investigated and confirmed by an allele-specific PCR, using a primer carrying the 5982G>A mutation at the 3'-end, in combination with the regular exon 30 primer set (Fig. 4). Clinical examination of the mother did not reveal any signs of CHARGE syndrome.

DISCUSSION

At the time of evaluation of our clinical data, *CHD7* sequencing had been performed in 107 index patients referred to our laboratory because of clinical features suggestive of CHARGE syndrome. Pathogenic mutations were identified in 69 patients (65%), including six patients that had previously tested negative.¹² All mutations except two were unique and most mutations had a severe effect on the *CHD7* protein, being either nonsense or frameshift mutations (70%).

From both our previous data and a recent report by Arrington et al. (2005) it is known that microdeletions of the chromosome 8q12.1 region, including the *CHD7* gene, may also result in CHARGE syndrome.¹²⁻¹⁵ We excluded the presence of such microdeletions in the patients without *CHD7* mutations by MLPA. From these results we conclude that whole gene deletions of the *CHD7* gene are not a frequent cause of CHARGE syndrome. Currently, we are extending our MLPA analyses in order to assess for the presence of small intragenic deletions.

In 20 out of 21 families a *de novo* occurrence of the *CHD7* mutation could be proven. In the mother of the sibs with the 5982G>A(W1994X) change, this mutation was present as a somatic mosaicism. It is likely that germline mosaicism exists as well. As a consequence prenatal diagnosis should be offered to all parents of children with an apparently *de novo* *CHD7* mutation.

Out of the 69 *CHD7* mutation-positive patients 45 index cases were selected for further clinical study together with two sibs, resulting in a cohort of 47 patients. Due to a short follow-up period, clinical information was limited in eleven patients, especially regarding hearing, growth and development. From the data presented in Table III and the detailed clinical description of our patients, it is clear that within the *CHD7* mutation-positive subset of CHARGE patients an extensive variability in clinical presentation exists, without any obvious genotype-phenotype correlation. This is best demonstrated in the two sib pairs. In the first sib pair, both twin girls had choanal atresia and a heart defect, but they were discordant for the coloboma and tracheo-esophageal fistula. The boys of the other sib pair were discordant for cleft lip/palate, heart defect, tracheo-esophageal fistula, coloboma and hearing loss. Missense mutations were found in three patients of the clinical study group, one of which was mildly mentally retarded. The other two had normal levels of intelligence. However, normal intelligence was also present in five patients with a nonsense mutation. Overall clinical comparison of these three patients with a missense mutation with the rest of the study group did not reveal any clear differences. However, it is still possible that less severe mutations (i.e. missense mutations) result in a less specific phenotype, not recognized as CHARGE syndrome. Hence, such patients may not be included in this study. On the other hand, patients with a *CHD7* deletion may be more severely affected than patients with a *CHD7* mutation, especially if multiple adjacent genes are deleted. Further studies are needed to explore this.

In Table IV the frequency of the main features of CHARGE syndrome in our group of *CHD7* mutation-positive patients is compared with data from the literature.

Table IV. The frequencies of characteristic CHARGE findings in a population of *CHD7*-positive patients compared to the literature

	This study N=47 (%)	Stromland et al. (2005) N=30 (%)	Issekutz et al. (2005) N=77 (%)	Tellier et al. (1998) N=47 (%)
Coloboma	33/47 70	90	77	79
Choanal atresia	17/47 36	35	64	57
Ear-anomaly/deafness	47/47 100	90	96	100
Cranial nerves	10/47* 21	32*	91	78
Genital hypoplasia (males)	17/22 77	67	65	53

Heart defects	31/47	66	52	84	85
Cleft lip/palate	17/47	36	19	18	17
TE fistula	8/47	17	16	19	15
Growth deficiency	21/32	66	48	58	75
Renal	9/47	19	12	36	19
Spine	9/47	19	27	9	13

* Data on facial nerve only

The distribution of features in the clinically diagnosed CHARGE syndrome patients as reviewed by Stromland et al. (2005), Issekutz et al. (2005) and Tellier et al. (1998) is comparable to the *CHD7* mutation-positive patients.^{3 10 16} This indicates that, within the patient group that fulfils the clinical diagnosis of CHARGE syndrome, there is not a specific subgroup that is more likely to have a *CHD7* mutation. None of the clinical features seems to be obligatory for a *CHD7* mutation, with the possible exception of vestibular anomalies. Several reports have stressed the high frequency and the high specificity of anomalies of the semicircular canals.^{1 2 17 18} This was also observed in our cohort of patients. All patients that were investigated by CT-scan or vestibular function tests had either abnormal function or an aplasia of the semicircular canals.

The effect of a *CHD7* mutation on a specific organ is variable and does not predict the consequences for other organ systems in which *CHD7* is expressed. For instance, a severe heart defect does not exclude normal intelligence (e.g. individual 43, Table III) and severe mental retardation does not have to be accompanied by severe defects in other organs (e.g. individual 8, Table III). This results in an enormous clinical variability, even within sib pairs. We carefully tested whether both sets of diagnostic criteria could be applied to our patients (see Table III).^{9 11} This was not possible in all cases, since for example CT-scanning of the temporal bones is required in order to apply the diagnostic criteria proposed by Verloes. For simplification we decided to use the 1998 Blake criteria as listed in Table I instead of the refined criteria adopted for different age groups.¹⁰ Both Blake and Verloes require that at least a coloboma or choanal atresia is present for the diagnosis CHARGE syndrome. Five patients in our study group (individuals 4-6-9-28-29 in Table III) neither had coloboma nor choanal atresia. Blake et al. (1998) argued that the choanae are usually patent when orofacial clefting is present and palatal clefting can be substituted for choanal atresia in the scoring criteria.⁹ As a consequence, only one patient (individual 28 in Table III) failed to fulfil the diagnostic criteria for CHARGE syndrome according to both Blake and Verloes. In 38 patients with features suggestive of CHARGE syndrome no *CHD7* mutation and/or deletion was identified. For 27 of these patients sufficient clinical data were available to apply the clinical diagnostic criteria. Only two of these 27 *CHD7* mutation-negative patients fulfilled the diagnostic criteria. In both patients aplasia of the semicircular canals was demonstrated. As a consequence the positive predictive value of the clinical diagnostic criteria is 37/39 (95%). This is substantiated by the fact that after improvement of the sequencing procedure (table I of supplemental data, *JMG* website) the mutation positive percentage in our first reported cohort reaches 95% (18/19).¹² In the context of the previously suggested genetic heterogeneity,¹⁹⁻²¹ this is an interesting observation that needs confirmation.

We would like to stress that CHARGE syndrome remains a clinical diagnosis. Although the high percentage of *CHD7* mutations in clinically diagnosed CHARGE syndrome patients indicates that *CHD7* is the major gene involved, this diagnosis cannot be rejected based on absence of a *CHD7* mutation. On the other hand, based on the clinical criteria alone, one *CHD7* positive patient would have been missed in our series.

In conclusion, we confirm that mostly unique *CHD7* mutations account for the majority of the cases with CHARGE syndrome, with a broad clinical variability and without an obvious genotype-phenotype correlation. In addition we provided evidence for germline mosaicism.

ACKNOWLEDGMENTS

We thank the patients and their parents for their participation and D. Wieczorek, T. Letteboer, J. Verheij, M. Baars, A. van Haeringen, J. Cobben, S. Maas, W. Kok, Y. Hilhorst-Hofstee, C. de Die-Smulders, M. Parisi, V. Der Kaloustian and B. Smyle for providing clinical data.

COMPETING INTERESTS

Non declared

LICENCE FOR PUBLICATION

The Corresponding Author has the right to grant on behalf of all authors and does grant on behalf of all authors, an exclusive licence (or non exclusive for government employees) on a worldwide basis to the BMJ Publishing Group Ltd to permit this article (if accepted) to be published in Journal of Medical Genetics and any other BMJ PGL products and sublicences such use and exploit all subsidiary rights, as set out in our licence (<http://JMG.bmjournals.com/misc/ifora/licenceform.shtml>).

INFORMED CONSENT

Informed consent was obtained of all patients that are represented by their photograph in this manuscript.

REFERENCES

- 1 **Morgan D**, Bailey M, Phelps P, Bellman S, Grace A, Wyse R. Ear-nose-throat abnormalities in the CHARGE association. *Arch Otolaryngol Head Neck Surg* 1993;**119**(1):49-54.
- 2 **Admiraal RJ**, Joosten FB, Huygen PL. Temporal bone CT findings in the CHARGE association. *Int J Pediatr Otorhinolaryngol* 1998;**45**(2):151-62.
- 3 **Tellier AL**, Cormier-Daire V, Abadie V, Amiel J, Sigaudy S, Bonnet D, de Lonlay-Debeney P, Morrissette-Durand MP, Hubert P, Michel JL, Jan D, Dollfus H, Baumann C, Labrune P, Lacombe D, Philip N, LeMerrer M, Briard ML, Munnich A, Lyonnet S. CHARGE syndrome: report of 47 cases and review. *Am J Med Genet* 1998;**76**(5):402-9.
- 4 **Hartshorne TS**, Grialou TL, Parker KR. Autistic-like behavior in CHARGE syndrome. *Am J Med Genet A* 2005;**133**(3):257-61.
- 5 **Smith IM**, Nichols SL, Issekutz K, Blake K. Behavioral profiles and symptoms of autism in CHARGE syndrome: Preliminary Canadian epidemiological data. *Am J Med Genet A* 2005;**133**(3):248-56.
- 6 **Hall BD**. Choanal atresia and associated multiple anomalies. *J Pediatr* 1979;**95**(3):395-8.
- 7 **Hittner HM**, Hirsch NJ, Kreh GM, Rudolph AJ. Colobomatous microphthalmia, heart disease, hearing loss, and mental retardation--a syndrome. *J Pediatr Ophthalmol Strabismus* 1979;**16**(2):122-8.
- 8 **Pagon RA**, Graham JM Jr, Zonana J, Yong SL. Coloboma, congenital heart disease, and choanal atresia with multiple anomalies: CHARGE association. *J Pediatr* 1981;**99**(2):223-7.
- 9 **Blake KD**, Davenport SL, Hall BD, Hefner MA, Pagon RA, Williams MS, Lin AE, Graham JM Jr. CHARGE association: an update and review for the primary pediatrician. *Clin Pediatr (Phila)* 1998;**37**(3):159-73.
- 10 **Issekutz KA**, Graham JM Jr, Prasad C, Smith IM, Blake KD. An epidemiological analysis of CHARGE syndrome: Preliminary results from a Canadian study. *Am J Med Genet A* 2005;**133**(3):309-17.
- 11 **Verloes A**. Updated diagnostic criteria for CHARGE syndrome: A proposal. *Am J Med Genet A* 2005;**133**(3):306-8.
- 12 **Vissers LE**, van Ravenswaaij CM, Admiraal R, Hurst JA, de Vries BB, Janssen IM, van der Vliet WA, Huys EH, de Jong PJ, Hamel BC, Schoenmakers EF, Brunner HG, Veltman JA, Geurts van Kessel AG. Mutations in a new member of the chromodomain gene family cause CHARGE syndrome. *Nat Genet* 2004;**36**(9):955-7.
- 13 **Delmas V**, Stokes DG, Perry RP. A mammalian DNA-binding protein that contains a chromodomain and an SNF2/SWI2-like helicase domain. *Proc Natl Acad Sci U S A* 1993;**90**(6):2414-8.
- 14 **Woodage T**, Basrai MA, Baxevanis AD, Hieter P, Collins FS. Characterization of the CHD family of proteins. *Proc Natl Acad Sci USA* 1997;**94**(21):11472-7.
- 15 **Arrington CB**, Cowley BC, Nightingale DR, Zhou H, Brothman AR, Viskochil DH. Interstitial deletion 8q11.2-q13 with congenital anomalies of CHARGE association. *Am J Med Genet* 2005;**133**(3):326-30.
- 16 **Stromland K**, Sjogreen L, Johansson M, Ekman Joelsson BM, Miller M, Danielsson S, Billstedt E, Gillberg C, Jacobsson C, Norinder JA, Granstrom G. CHARGE association in Sweden: Malformations and functional deficits. *Am J Med Genet A* 2005;**133**(3):331-9.
- 17 **Amiel J**, Attiee-Bitach T, Marianowski R, Cormier-Daire V, Abadie V, Bonnet D, Gonzales M, Chemouny S, Brunelle F, Munnich A, Manach Y, Lyonnet S. Temporal bone anomaly proposed as a major criteria for diagnosis of CHARGE syndrome. *Am J Med Genet* 2001;**99**(2):124-7.
- 18 **Satar B**, Mukherji SK, Telian SA. Congenital aplasia of the semicircular canals. *Otol Neurotol* 2003;**24**(3):437-46.
- 19 **Clementi M**, Tenconi R, Turolla L, Silvan C, Bortotto L, Artifoni L. Apparent

- CHARGE association and chromosome anomaly: chance or contiguous gene syndrome. *Am J Med Genet* 1991;**41**(2):246-50.
- 20 **North KN**, Wu BL, Cao BN, Whiteman DA, Korf BR. CHARGE association in a child with de novo inverted duplication (14)(q22-->q24.3). *Am J Med Genet* 1995;**57**(4):610-4.
- 21 **Martin DM**, Sheldon S, Gorski JL. CHARGE association with choanal atresia and inner ear hypoplasia in a child with a de novo chromosome translocation t(2;7)(p14;q21.11). *Am J Med Genet* 2001;**99**(2):115-9.

FIGURE LEGENDS

Legend to Figure 1

Distribution of *CHD7* mutations identified in the 69 CHARGE syndrome patients. Coding exons are indicated in black bars, whereas the non-coding sequences are indicated in gray. Colored bars represent chromodomains (red), SNF2 domain (green) and Helicase domain (yellow). Mutations are schematically shown above the exons in which they are located. Nonsense mutations are represented by ● (n=31), missense mutations by ■ (n=8), frameshift mutations (n=17) by ◆ and splice site mutations by ▲ (n=13) respectively.

Legend to Figure 2

Patient 28, who is *CHD7* mutation positive but does not fulfil the diagnostic criteria (see text). (Written consent was obtained for publication of this picture)

Legend to Figure 3

a), twin 1 of sib pair 1 (patient 27 in table III), who died shortly after birth; b) twin 2 of sib pair 1 (patient 26 in table III) at the age of 7 years; c) sib 1 and d) sib 2 of sib pair 2, both at the age of 2 years (patients 29 and 30 in table III). (Written consent was obtained for publication of these pictures)

Legend to Figure 4

Results of allele-specific PCR for 5982G>A mutation in the family with two affected boys (a). DNAs of the indicated family members and an unrelated unaffected control (IV) were subjected to a multiplex PCR using a mutation-specific primer (5982G>A, lower band) and the regular primer set for exon 30 (upper band) (b). The mutation found in the boys (I) was also present in the mother (II). The different relative amounts of the fragments of individual I and II, might reflect the presumed mosaicism in the mother.

Figure 1






Figure 2

Figure 3a



Figure 3b



Figure 3c



Figure 3d





Figure 4

a.



b.




Table I Primers for sequence analysis of CDH7 gene.

Note: All forward primers have an M13 universal tail, all reverse primers have an M13-reverse universal tail to allow easy sequence analysis.

The primers marked with an asterisk (*) are different from the primers used in our original article on the identification of the *CHD7* gene (*Nat Genet* 2004;36(9):955-7).

Primer	Exon/Fragment	F/R	Sequence
6386	02a*	F	tgtaaaacgacggccagttattataaaatacagctcagcgaggggagc
6465	02a*	R	caggaaacagctatgaccgtagggggctcgtatctggt
6466	02b*	F	tgtaaaacgacggccagtcggtcagatgggtgtctacc
6467	02b*	R	caggaaacagctatgaccaagttccctgaggctgttgaa
6468	02c*	F	tgtaaaacgacggccagtaccacagtggtcacttaac
6387	02c*	R	caggaaacagctatgaccggcactcataatgaaaaagctttcatagag
6388	03*	F	tgtaaaacgacggccagtgatgatagattctacaactgtatagtgg
6389	03	R	caggaaacagctatgaccatttcataggctgtaactatttggcgc
6390	04	F	tgtaaaacgacggccagtgcattttgcacaaatgtactatgtgtgg
6391	04	R	caggaaacagctatgaccggggaggctctgtgtacttcatacatgg
6392	05	F	tgtaaaacgacggccagtcggcctcccaaagtgctgggattacagg
6469	05*	R	caggaaacagctatgaccactgcacttctcagattgcagcaatg
6470	06	F	tgtaaaacgacggccagtcagtgacttaaaaggtgtggaggtgg
6393	06	R	caggaaacagctatgacctagacaggattagaaatattactggaag
6394	07	F	tgtaaaacgacggccagtcctcaaatggggcaggttgtgtg
6395	07	R	caggaaacagctatgaccatacacagaagttagtcaactttacacc
6396	08	F	tgtaaaacgacggccagtgctcagcagccttaatgggtaattaagc
6471	08	R	caggaaacagctatgaccgtaccaatagcaagttgacagcac
6472	09	F	tgtaaaacgacggccagtcctcaaatgtaagttttatattgcttgacc
6397	09	R	caggaaacagctatgaccctcaagtgctcacagtcctcaaggctctg
6398	10-11	F	tgtaaaacgacggccagtatgtgatgtatgtggctcaaatgaatcc
6473	10-11	R	caggaaacagctatgaccgtctttttattatttcaataactaaagg
6474	12	F	tgtaaaacgacggccagtttgggtatgcatttgggtacaatgg
6475	12*	R	caggaaacagctatgacccttccaagtcaccaagac
6476	13	F	tgtaaaacgacggccagttataactgaaaacagaatgtatgtcacc
6399	13	R	caggaaacagctatgaccccagagaatatatcacaatattgacaagc
6400	14	F	tgtaaaacgacggccagtcctgattctatactttgcatagggtag
6477	14	R	caggaaacagctatgaccggtgggaaaaactgtaataatacacac
6478	15	F	tgtaaaacgacggccagttggatgtttaatgaatgagataatcctg
6401	15	R	caggaaacagctatgaccacctgctgagtgagcaggagacagg
6402	16	F	tgtaaaacgacggccagttattctgttcataagcaggagtttggg
6480	16	R	caggaaacagctatgaccttttaggtggactgctggacccc
6481	17*	F	tgtaaaacgacggccagtcgccaataaacctatttgc
6482	17	R	caggaaacagctatgaccgcaacattaatttattgggtctgactgg
6483	18-19	F	tgtaaaacgacggccagtcctcaaatccatcacattggaatgagg
6403	18-19	R	caggaaacagctatgaccattattccaatgcatcttgaagcac
6404	20-21	F	tgtaaaacgacggccagtagtgagggaatgcacaatatcggagca
6484	20-21	R	caggaaacagctatgaccacaaatagttttctctaccagttgg
6485	22-23	F	tgtaaaacgacggccagtcgctggtacctgacttaaaagtaagcc
6406	22-23	R	caggaaacagctatgaccgcatctgcagacgtgaagggccggctgc
6405	24-25	F	tgtaaaacgacggccagtgccaggatgatgatgaacagcagcagc
6486	24-25	R	caggaaacagctatgacctatgactttggaaaagagatgtaaagag
6487	26*	F	tgtaaaacgacggccagtggtgtgtggcagtgctgtgatttggca
6488	26	R	caaaaaaacactataaccnaaatnaatnaaaaaaanaataaatct
6489	27-28	F	tgtaaaacgacggccagtagattattactcttctaccacccc
6407	27-28	R	caggaaacagctatgaccacagtgacaatgactgctcagtcac

6490	29	F	tgtaaacgacggccagtgactgagatgcccttcccactgtc
6491	29	R	caggaacagctatgacccttctccccagattatacatgg
6492	30	F	tgtaaacgacggccagtcctatgataaagctggggggaagaag
6493	30	R	caggaacagctatgaccctcgtcttttqtatagaactttgtt
6494	31	F	tgtaaacgacggccagtaacaaagttctatacaaaaagacgag
6495	31	R	caggaacagctatgaccgcaggtgccgtgctgccagaaagcaacg
6496	32	F	tgtaaacgacggccagtcctatgtagtaggtactcaataaaatggagc
6409	32	R	caggaacagctatgaccaagctagaagattcctctaccctaac
6497	33	F	tgtaaacgacggccagtcatttatgctcttttgcactttgatgg
6498	33	R	caggaacagctatgaccgggctggcttttagaataaggaaca
6499	34	F	tgtaaacgacggccagttgttccttattctaaaagccagccc
6411	34	R	caggaacagctatgaccggcttcatacaatgctgctgagagaaac
6412	35	F	tgtaaacgacggccagttcccaacaactagacattgtttctag
6500	35	R	caggaacagctatgaccctgtcagggatttctatgttqtaagg
6501	36-37	F	tgtaaacgacggccagtttgaagatgatctgacagttctcttgg
6413	36-37	R	caggaacagctatgaccgatgtattatgtcaattctttaaagctaag
6502	38a	F	tgtaaacgacggccagttccaccacagaggctcacattgagatc
6503	38a*	R	caggaacagctatgacctcgtcttcattctcatttcc
6504	38b*	F	tgtaaacgacggccagtaggagaaccggaagacag
6505	38b*	R	caggaacagctatgaccgcactgcacaataacttaatgac

Table II Conditions for sequence analysis of CDH7 gene.

Fragment / Exon	Forward primer		Reverse primer		Fragment length (bp)	PCR conditions							
	Number	Conc. (ng/μl)	Number	Conc. (ng/μl)		Buffer (Applied Biosystems)	Denaturation		Annealing		Elongation		Cycli
							Temp. (°C)	time	Temp. (°C)	time	Temp. (°C)	time	
2a	6386	100	6465	100	680	AB gold + 2,0 mM MgCl ₂	94	30"	55	30"	72	1'	35
2b	6466	100	6467	100	750	AB gold + 2,0 mM MgCl ₂	94	30"	55	30"	72	1'	35
2c	6468	100	6387	100	622	AB gold + 2,0 mM MgCl ₂	94	30"	55	30"	72	1'	35
3	6388	100	6389	100	659	AB gold + 2,0 mM MgCl ₂	94	30"	55	30"	72	1'	35
4	6390	100	6391	100	485	AB gold + 2,0 mM MgCl ₂	94	30"	55	30"	72	1'	35
5	6469	100	6392	100	332	AB gold + 1,5 mM MgCl ₂ + DMSO	94	30"	55	30"	72	1'	35
6	6470	100	6393	100	408	AB gold + 2,0 mM MgCl ₂	94	30"	55	30"	72	1'	35
7	6394	100	6395	100	568	AB gold + 1,5 mM MgCl ₂ + DMSO	94	30"	55	30"	72	1'	35
8	6396	100	6471	100	374	AB gold + 2,0 mM MgCl ₂	94	30"	55	30"	72	1'	35
9	6472	100	6397	100	320	AB gold + 2,0 mM MgCl ₂	94	30"	55	30"	72	1'	35
10-11	6398	100	6473	100	493	AB gold + 2,0 mM MgCl ₂	94	30"	55	30"	72	1'	35
12	6474	100	6475	100	367	AB gold + 2,0 mM MgCl ₂	94	30"	55	30"	72	1'	35
13	6476	100	6399	100	452	AB gold + 2,0 mM MgCl ₂	94	30"	55	30"	72	1'	35
14	6400	100	6477	100	382	AB gold + 2,0 mM MgCl ₂	94	30"	55	30"	72	1'	35
15	6478	100	6401	100	483	AB gold + 2,0 mM MgCl ₂	94	30"	55	30"	72	1'	35
16	6402	100	6480	100	418	AB gold + 2,0	94	30"	55	30"	72	1'	35

						mM MgCl ₂							
17	6481	100	6482	100	376	AB gold + 2,0 mM MgCl ₂	94	30"	55	30"	72	1'	35
18-19	6483	100	6403	100	793	AB gold + 2,0 mM MgCl ₂	94	30"	55	30"	72	1'	35
20-21	6404	100	8484	100	598	AB gold + 2,0 mM MgCl ₂	94	30"	55	30"	72	1'	35
22-23	6485	100	6406	100	713	AB gold + 2,0 mM MgCl ₂	94	30"	55	30"	72	1'	35
24-25	6405	100	6486	100	825	AB gold + 2,0 mM MgCl ₂	94	30"	55	30"	72	1'	35
26	6487	100	6488	100	562	AB gold + 2,0 mM MgCl ₂	94	30"	55	30"	72	1'	35
27-28	6489	100	6407	100	437	AB gold + 1,5 mM MgCl ₂	94	30"	55	30"	72	1'	35
29	6490	100	6491	100	534	AB gold + 2,0 mM MgCl ₂	94	30"	55	30"	72	1'	35
30	6492	100	6493	100	344	AB gold + 1,5 mM MgCl ₂	94	30"	55	30"	72	1'	35
31	6494	100	6495	100	862	AB gold + 2,0 mM MgCl ₂	94	30"	55	30"	72	1'	35
32	6496	100	6409	100	355	AB gold + 2,0 mM MgCl ₂	94	30"	55	30"	72	1'	35
33	6497	100	6498	100	509	AB gold + 2,0 mM MgCl ₂	94	30"	55	30"	72	1'	35
34	6499	100	6411	100	606	AB gold + 2,0 mM MgCl ₂	94	30"	55	30"	72	1'	35
35	6412	100	6500	100	451	AB gold + 2,0 mM MgCl ₂	94	30"	55	30"	72	1'	35
36-37	6501	100	6413	100	678	AB gold + 2,0 mM MgCl ₂	94	30"	55	30"	72	1'	35
38a	6502	100	6503	100	562	AB gold + 2,0 mM MgCl ₂	94	30"	55	30"	72	1'	35
38b	6504	100	6505	100	685	AB gold + 2,0 mM MgCl ₂	94	30"	55	30"	72	1'	35

Table 3

EXON	MLPA-primer
CHARGE_2af	gggttcctaagggttggac ctcagtgaagtgaagcacag
CHARGE_2bf	gggttcctaagggttggac tccaccatcaccccagaacac
CHARGE_3f	gggttcctaagggttggac ggagcccttctagagaaaccag
CHARGE_4f	gggttcctaagggttggac tagacaaaacccccaccatctcctc
CHARGE_5f	gggttcctaagggttggac aggaggcagatgatgcagatgctgctgggag
CHARGE_6f	gggttcctaagggttggac aggtgagtgccattggagccattaaaatctgt
CHARGE_7f	gggttcctaagggttggac ctcccagcacacggtgtgctcatcttacgtgaa
CHARGE_8f	gggttcctaagggttggac tctctctctctctctctctctgtaacaggttaagag
CHARGE_9f	gggttcctaagggttggac tatgtggaggtgaccggataatggacttgcacgtagca
CHARGE_10_11f	gggttcctaagggttggac gagcggaggcaggacatagatcaagcaagatcgaggagttg
CHARGE_33f	gggttcctaagggttggac ggcgccagatgtttgattcc
CHARGE_34f	gggttcctaagggttggac aatggaactgctccaagcaggcc
CHARGE_35f	gggttcctaagggttggac attcctctcccggacagctgga
CHARGE_36f	gggttcctaagggttggac gggtgctgttgcaataaacgaaatgg
CHARGE_37f	gggttcctaagggttggac tcaaatgcctctaccagatgggtggagct
CHARGE_38f	gggttcctaagggttggac agatgctgttctggctgctgactctgcaatgg
CHARGE_2ar	gcaagctcctgagctgtggtt ctagattggatcttgcctggcac
CHARGE_2br	cccgcagaaagtgcctgtgcat ctagattggatcttgcctggcac
CHARGE_3r	tgccggatatgactcaggtagtggt ctagattggatcttgcctggcac
CHARGE_4r	ctcctgaagaagatgaggaccaggtgtt ctagattggatcttgcctggcac
CHARGE_5r	gattccccctccaacacctcccagtcagaat ctagattggatcttgcctggcac
CHARGE_6r	gaggtgatgtgactcttacaggattgttgct ctagattggatcttgcctggcac
CHARGE_7r	ggtccttgctgctctttcagtcctat ctagattggatcttgcctggcac
CHARGE_8r	agctggaggcttaggacagaaggcaa ctagattggatcttgcctggcac
CHARGE_9r	cagatgaccggggagaggtaacaggagatcattgt ctagattggatcttgcctggcac
CHARGE_10_11r	agaaactaatgtccaggagccggaacagagcgtgtgta ctagattggatcttgcctggcac
CHARGE_33r	aaggctcatcccaggttacat ctagattggatcttgcctggcac
CHARGE_34r	ttcgcgcacaccacaaggcatt ctagattggatcttgcctggcac
CHARGE_35r	cccagacacaggatccctgttat ctagattggatcttgcctggcac
CHARGE_36r	gaagaaggtaaacgctgggaaagggaatt ctagattggatcttgcctggcac
CHARGE_37r	atggcgctccaatgaaggatctaccaggt ctagattggatcttgcctggcac
CHARGE_38r	atctgttgctgctactgccccggctggatt ctagattggatcttgcctggcac