OPEN ACCESS

Original research

De novo mutations of *SCN1A* are responsible for arthrogryposis broadening the *SCN1A*-related phenotypes

Dana Jaber,¹ Cyril Gitiaux,² Sophie Blesson,³ Florent Marguet,^{4,5} David Buard,¹ Maritzaida Varela Salgado,¹ Anna Kaminska,⁶ Julien Saada,⁷ Catherine Fallet-Bianco,⁸ Jelena Martinovic,⁹ Annie Laquerriere,^{4,5} Judith Melki ^{1,10}

For numbered affiliations see end of article.

Correspondence to

Professor Judith Melki, INSERM, Paris 94276, France; judith.melki@inserm.fr

Received 8 May 2020

Revised 2 August 2020

Accepted 6 August 2020

ABSTRACT

Background Arthrogryposis multiplex congenita (AMC) is the direct consequence of reduced fetal movements. AMC includes a large spectrum of diseases which result from variants in genes encoding components required for the formation or the function of the neuromuscular system. AMC may also result from central nervous involvement. *SCN1A* encodes Nav1.1, a critical component of voltage-dependent sodium channels which underlie action potential generation and propagation. Variants of *SCN1A* are known to be responsible for Dravet syndrome, a severe early-onset epileptic encephalopathy. We report pathogenic heterozygous missense de novo variants in *SCN1A* in three unrelated individuals with AMC.

Methods Whole-exome sequencing was performed from DNA of the index case of AMC families. Heterozygous missense variants in *SCN1A* (p.Leu893Phe, p.Ala989Thr, p.Ile236Thr) were identified in three patients. Sanger sequencing confirmed the variants and showed that they occurred de novo.

Results AMC was diagnosed from the second trimester of pregnancy in the three patients. One of them developed drug-resistant epileptic seizures from birth. We showed that *SCN1A* is expressed in both brain and spinal cord but not in skeletal muscle during human development. The lack of motor denervation as established by electromyographic studies or pathological examination of the spinal cord or skeletal muscle in the affected individuals suggests that AMC is caused by brain involvement.

Conclusion We show for the first time that *SCN1A* variants are responsible for early-onset motor defect leading to AMC indicating a critical role of *SCN1A* in prenatal motor development and broadening the phenotypic spectrum of variants in *SCN1A*.

Considerable genetic heterogeneity has been reported in AMC. AMC includes a large spectrum of diseases which result from variants in genes encoding components required for the formation or the function of neuromuscular junctions, skeletal muscle, survival of motor neurons, myelination of peripheral nerves or the function of connective tissues.³ AMC may also result from central nervous involvement including channelopathies linked to variants of *NALCN* (MIM: 611549)⁴ or *CACNA1E* (MIM: 601013).⁵

The identification of the pathogenic variant in AMC is essential to provide information regarding prognosis and comorbidities and informs accurate genetic counselling. Here, we further explored genetic alterations in a group of genetically undiagnosed AMC through whole-exome sequencing (WES) and identified heterozygous de novo missense variants in *SCN1A* in three unrelated patients. *SCN1A* (MIM: 182389) encodes a critical component of voltage-dependent sodium channels. Variants of this gene most frequently present as Dravet syndrome (EIEE6, [MIM: 607208]), an early-onset epileptic encephalopathy.⁶ We present the clinical data for these three patients and describe the phenotypic spectrum associated with this new recurrent genetic disorder.

METHODS

The parents of all affected individuals provided written informed consents for pathological examinations and genetic analyses of their affected children or fetuses and themselves in accordance with the ethical standards of our institutional review boards.

Whole-exome sequencing

WES was performed from DNA of the index case in the AMC families using a completed Twist Bioscience Human Core Exome (Consensus CDS) kit for library preparation and exome enrichment (Integrage). Sequencing was performed on a Genome Analyzer HiSeq4000 Illumina instrument in paired-end mode with a read length of 2×75 bp (Integrage). Reads were aligned to the human reference genome sequence (UCSC hg19, NCBI build 37.3) via the BWA program.⁷ Variants were selected using the SAMtools⁸ then annotated using Annovar software.⁹ Variants in coding regions

INTRODUCTION

Arthrogryposis multiplex congenita (AMC) has an overall incidence of 1 in 3000 to 5000.^{1,2} AMC is the direct consequence of reduced fetal movements which may lead, in addition to AMC, to pterygia, pulmonary hypoplasia, diaphragmatic defect or cleft palate. Some non-genetic factors may cause AMC such as extrinsic limitation of fetal movements, bacterial or viral infections, or maternal autoimmune myasthenia.

© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Jaber D, Gitiaux C, Blesson S, et al. *J Med Genet* Epub ahead of print: [please include Day Month Year]. doi:10.1136/jmedgenet-2020-107166

(including non-synonymous and nonsense variants), intron-exon junctions or short coding insertions or deletions were selected when the minor allele frequency was less than 0.005. Prediction of pathogenicity of missense variants was performed using Polyphen-2¹⁰ and SIFT software.¹¹

Sanger sequencing

Variants identified through WES were validated by Sanger sequencing. The primers used for the detection of the *SCN1A* c.2679A>C variant are 5'-CTATCCACTCCCCACACAGC-3' and 5'-GAGACGGTTAGGGCAGATCA-3'; for c.2965G>A: 5'-AGCTACTCCTTTGTGCATCCT-3' and 5'-CAGTCACA AATTCAGATCACCCA-3' and for c.707T>C: 5'-CTGCAG-CCCAATTAGAGCAA-3' and 5'-CTTCTCCACTAGC-GTTGCAA-3'. PCR amplification was carried out as previously described.¹² PCR products were purified then sequenced using the forward or reverse primers (Eurofins Genomics). The obtained DNA sequences were compared with published sequences (BLAST, NCBI). Sanger sequencing was also performed to establish the genotype of each family member and to analyse the segregation of the variant within each family.

Real-time PCR amplification of genomic DNA

Real-time PCR amplification was conducted using genomic DNA on a 7300 Real-Time PCR system (Applied Biosystems) using Power SYBR Green PCR Master Mix (Applied Biosystems).¹² Genomic deletion was defined when the ratio of tested DNA to control DNA was equal to or less than 0.5. Real-time PCR amplification of each sample was performed in duplicate using primers within *SCN1A* exons 1, 10, 14, 15 and 16 (data available on request). Albumin gene was used as internal control (data available on request).

RT-PCR amplification

Total RNAs were extracted from samples of frontal cortex, spinal cord or skeletal muscle by using TRI Reagent LS method (Sigma-Aldrich). RNA (1.5 µg) was used to synthesise cDNA by using random hexamers following the manufacturer's instructions (SuperScript III reverse Transcriptase; Invitrogen) in a final volume of 20 µL. PCR amplification was carried out using primers chosen in exons of *SCN1A* (5'-GAAGAACA GCCCGTAGTGGA-3'; 5'-TTCAAATGCCAGAGCACCA-3') as previously described.¹² RT-PCR products were separated by 2% agarose gel electrophoresis and labelled with ethidium bromide. As internal control for PCR amplification, β-actin cDNA was co-amplified (5'-GCCAGGTCATCACCATTGG-3'; 5'-AGGACTCCATGCCAGGAA-3').

RESULTS

Identification of de novo heterozygous missense *SCN1A* variants in arthrogryposis

In family 1, the affected child was born to non-consanguineous healthy parents (figure 1). During pregnancy and based on ultrasound examination, the male fetus displayed from 23 weeks of gestation (w.g.) bilateral flexion of both hands, a hyperextension of knees and reduced swallowing. These data were confirmed at the third trimester ultrasound examination. The child was born at 38 w.g. and displayed microretrognathism, distal joint contractures of both lower and upper limbs, severe hypotonia and respiratory distress (figure 2). From postnatal day 2, he developed drug-resistant epileptic seizures. Six video EEGs were performed between the age of 2 and 15 days of life, showing dysmature aspect with high amplitude temporal and central delta brushes and discontinuous (up to 15 s of inter-burst interval) and intermittent asynchronous activity in quiet sleep

Figure 1 De novo variants in *SCN1A* in three arthrogryposis multiplex congenita families and transcript analysis. (A) Pedigrees with Sanger sequencing results for families 1, 2 and 3 are shown. Arrows indicate mutant nucleotide positions. The affected individuals carry de novo heterozygous *SCN1A* variants. The nucleotide and amino acid changes based on NM_001165963.2 and NP_001159435.1 reference sequences respectively are indicated. Open symbols: unaffected; filled symbols: affected. (B) RT-PCR amplification was performed using *SCN1A* specific primers on RNAs extracted from the brain motor cortex (BR), spinal cord (SC) and skeletal muscle (SM) of human fetuses at 27 or 22 w.g. When compared with the β-actin control, *SCN1A* is expressed in brain and spinal cord but not in skeletal muscle. MW, molecular weight.

Figure 2 Arthrogryposis multiplex congenita phenotype of the affected individuals. (A, A') The affected child of family 1 displayed distal joint contractures including camptodactyly and dorsal flexion of feet. (B) Postmortem examination of the affected fetus of family 3 confirmed bilateral camptodactyly, hyperextension of knees and hallux valgus of feet.

without typical aspect of suppression burst. Frequent rolandic and temporal sharp waves and bi-occipital and central seizures were recorded with subtle clinical features (figure 3). Brain MRI, metabolic investigation and array CGH were normal. Electromyography (EMG) and nerve conduction velocity studies revealed no evidence for peripheral neuropathy nor motor neuron involvement. Morphological analysis of muscle biopsy did not reveal any sign suggesting a congenital myopathy or muscle denervation. The patient was sequentially treated by the following drugs without efficacy on seizure control: phenobarbital, phenytoin, pyridoxine and clonazepam. The newborn died at 21 days from complete ventilator dependency and intractable epilepsy. WES was performed from DNA of the index case. A heterozygous variant was identified in *SCN1A* in the affected individual of family 1 (figure 1). It is a missense variant (NM_001165963.2:c.2679A>C) predicted to be pathogenic

Figure 3 Inter-ictal and ictal EEG at day 2 of patient 1. (A) Active sleep: frontal sharp transients and continuous slow-wave activity mixed with abundant rapid rhythms predominating on central regions (Delta brushes). (B) Quiet sleep with discontinuous activity and bursts composed of high-amplitude slow waves mixed with rapid rhythms and sharp waves. (C) Ictal EEG with seizures beginning on right occipital region, spreading to left occipital and bi-central regions.

with a high score (p.Leu893Phe; Polyphen-2 score of 1). This variant is not annotated in dbSNP153, 1000 Genomes, ExAC nor TOPMED databases. The variant was confirmed by Sanger sequencing using primers flanking the variant in the affected child. Analysis of both parents showed that the variant had occurred de novo (figure 1).

Analysis of WES previously performed on the DNA samples from additional affected individuals with AMC and/or reduced fetal mobility of unknown origin (n=147) was revised and allowed the identification of predicted pathogenic heterozygous variants in *SCN1A* in two additional individuals. In family 2, based on ultrasound examination at 25 w.g., the male fetus displayed a lack of fetal movements associated with arthrogryposis of the upper limbs and microretrognathism. The pregnancy was terminated at 25 w.g. at the request of the parents. Postmortem examination confirmed micrognathia without cleft palate, generalised oedema, very short neck, hyperextension of both legs in front of the trunk, dorsal flexion of feet, bilateral camptodactyly with flexion of elbows, right diaphragmatic hernia and marked dorsolumbar scoliosis. Neuropathological examination did not reveal any brain or spinal cord abnormalities. In the affected fetus of family 2, a heterozygous missense variant (NM_001165963.2:c.2965G>A) predicted to be pathogenic with a high score (p.Ala989Thr; Polyphen-2 score of 1) was identified. This variant is not annotated in databases and was confirmed by Sanger sequencing using primers flanking the variant in the affected fetus. Analysis of both parents revealed that the variant had occurred de novo (figure 1).

In family 3 (figure 1), based on ultrasound examination performed at 21 w.g., the male fetus displayed bilateral camptodactyly, hyperextension of knees and hallux valgus of feet. The pregnancy was terminated at 24 w.g. at the request of the parents. Postmortem examination confirmed the ultrasound findings (figure 2). Neuropathological examination did not reveal brain nor spinal cord abnormalities. A heterozygous missense variant (NM_001165963.2:c.707T>C) predicted to be pathogenic with a high score (p.Ile236Thr; Polyphen-2 score of 0.99) was identified in the affected fetus. This variant is annotated in dbSNP153 (rs886039464) database with unknown frequency. This variant was confirmed by Sanger sequencing using primers flanking the variant in the affected fetus. Analysis of both parents showed that the variant had occurred de novo (figure 1).

Looking at the position of variants of *SCN1A* in the three affected individuals did not reveal a specific domain of the Nav1.1 protein when compared with reported de novo pathogenic variants of *SCN1A* in Dravet syndrome (figure 4, Clinvar, NCBI). Since the disease-causing mechanism is a loss-of-function in a single copy of the *SCN1A* gene in Dravet syndrome, allelic variants of *SCN1A* could be responsible for more severe phenotypes. Through re-analysis of WES data and real-time PCR performed on DNA of affected individuals using primers located in exons 1, 10, 14, 15 and 16 of *SCN1A*, no additional variant including insertion/deletion in *SCN1A* was detected in the three affected individuals.

Mice with heterozygous deletion of *Scn1a* (*Scn1a*^{+/-}) model a number of Dravet syndrome features, including spontaneous seizures and premature lethality. Phenotypic severity in *Scn1a*^{+/-} mice strongly depends on strain background.¹³ On the 129S6/SvEvTac strain, *Scn1a*^{+/-} mice exhibit no overt phenotype, while on the (C57BL/6J×129S6/SvEvTac)F1 strain, *Scn1a*^{+/-} mice exhibit spontaneous seizures and early lethality suggesting modifier loci.¹³ Using genetic mapping on backcrosses, Miller *et al*¹³ identified several candidate loci that influence premature lethality of *Scn1a*^{+/-} mice. Differential

Figure 4 Protein structure of SCN1A and distribution of disease-causing missense variants found in patients with arthrogryposis multiplex congenita (AMC) compared with those found in Dravet syndrome. SCN1A is a monomer and consists of four homologous domains (DI–DIV). Each domain has six transmembrane and an intramembrane domains (the central pore). The positions of variants that cause AMC are indicated by vertical arrows. The percentage indicates the proportion of reported de novo pathogenic variants relative to the size of each domain and found in Dravet syndrome.

expression of genes within these candidate loci have been reported including *RELN*, *LGI2*, *GABRA2*, *MAPK10*, *ATP1A3*, *SLC7A10*, *GABRG3*, *KCNJ11*, *GABRB3*, *CLCN3*, *CACNA1A*, *GABRA6* and *GABRB2*.¹³ These genes might be regarded as modifiers in the affected AMC individuals reported here. However, we did not identify any predicted pathogenic variant in these candidate genes.

We also looked at genes encoding other components of axon initial segment (AIS)¹⁴ under the hypothesis of a loss of compensatory mechanism of genes encoding other components of AIS such as Nav1.2, Nav1.6 to 1.9, or *NALCN* or *CACNA1E*, two genes encoding cation channels and reported as responsible for AMC resulting from central nervous involvement.^{4,5} No predicted pathogenic variants in these genes were identified in the affected AMC individuals. Finally, we also tested the hypothesis of modifying genes unrelated to these genes. To this end, reanalysis of WES of the three affected individuals was performed. No predicted pathogenic variant in genes shared by the affected individuals except for *SCN1A* was identified. A larger number of affected individuals is likely required to isolate relevant modifier signal.

Arthrogryposis caused by *SCN1A* mutation results from brain involvement

SCN1A encodes Nav1.1 belonging to voltage-dependent sodium channels that regulate sodium exchange in excitable tissues including brain, spinal cord and muscle.¹⁵ Duflocq *et al*¹⁶ showed that Nav1.1 expression is concentrated in the proximal AIS and nodes of Ranvier throughout the mouse spinal cord and in many brain regions including cerebral cortex, basal ganglia and cerebellum. The AIS is a specialised structure in neurons that resides between axonal and somatodendritic domains. It serves two functions: (1) to integrate synaptic inputs and generate action potentials and (2) to maintain neuron polarity.¹⁷ The myelin sheath produced by Schwann cells and oligodendrocytes attaches to the axon at both sides of the Nodes of Ranvier creating specialised and regularly spaced domains. The propagation of action potentials in myelinated axons requires the precise localisation of Na⁺ channels at the nodes of Ranvier. These specialised domains underlie saltatory conduction of action potentials along myelinated axons, an essential process for neuronal function.¹⁴ To determine whether the AMC phenotype resulted from brain, spinal cord or skeletal muscle defect, *SCN1A* RNA expression analysis was performed from brain motor cortex, spinal cord or

skeletal muscle samples from 22 w.g. and 27 w.g. human fetuses affected from unrelated lethal disorders. RT-PCR analysis revealed an expression of *SCN1A* in brain and spinal cord but not skeletal muscle when compared with β -actin expression at w.g. similar to the age of AMC detection in the affected fetuses (figure 1). The RNA expression of *SCN1A* was similar to that reported in GTEx Portal (GTEx Analysis Release V8).

Gitiaux *et al*¹⁸ reported several cases with Dravet syndrome and EMG recordings showed features of chronic denervation. However, in case 1, EMG and nerve conduction velocity did not reveal any reduction of nerve conduction velocity nor features consistent with chronic denervation. Consistently, Spagnoli *et al*¹⁹ reported a case with severe, pharmacoresistant epileptic encephalopathy with multiple seizure types including tonic spasms associated with severe motor defect including contractures and talipes. Brain MRI and EMG were also normal.

These data strongly suggest that the AMC reported in these three individuals is likely the consequence of brain involvement. In addition, the normal nerve conduction velocity performed in case 1 indicates the absence of node of Ranvier defect of the peripheral nerve in contrast to AMC linked to variants in either *GLDN* (gliomedin, [MIM: 608603]) or *CNTNAP1* (contactin-associated protein 1, [MIM: 602346]) encoding essential components of the nodes of Ranvier and paranodes, respectively, and showing marked defect of peripheral nerve.^{20,21}

The expression pattern of *SCN1A* as well as the phenotype of the affected fetuses indicate that *SCN1A* has a critical role during prenatal motor development likely linked to the role of Nav1.1 in motor cortex.

DISCUSSION

Variants in Nav1.1 cause several related disorders including (1) Dravet syndrome, an early-onset epileptic encephalopathy (EIEE6, [MIM: 607208])⁶, (2) generalised epilepsy with febrile seizures plus (GEFSP2, [MIM: 604403])²² and (3) familial hemiplegic migraine as the main phenotypes (FHM3, [MIM: 609634]).²³ In Dravet syndrome which represents the most severe disease phenotype, patients often present with drug-resistant epilepsy beginning in the first year of life. In addition, patients with Dravet syndrome may also suffer from various symptoms including intellectual disability, and behavioural and motor impairment. The incidence of Dravet syndrome is 1 of 15 500.²⁴ Surprisingly, extensive neuropathological examination has not shown any consistent cerebral or spinal cord structural abnormalities, cell loss or neurodegenerative process in this condition.²⁵

Our approach allowed the identification of de novo missense *SCN1A* variants in three unrelated individuals with AMC. This conclusion is strongly supported by multiple lines of evidence: the variants are de novo and are absent from various control databases; the variants occur at residues that exhibit a high level of evolutionary conservation. In one patient, the phenotype was characterised by the association of AMC with severe and early-onset drug-resistant epileptic seizures and abnormal EEG background activity. The abnormal motor phenotype was detected from the second trimester of pregnancy and characterised by AMC in the three affected individuals. *SCN1A* expression during human development, EMG and morphological analysis of the affected individuals strongly suggest that AMC is linked to *SCN1A* defect in motor cortex.

The expression pattern of *SCN1A* in the central nervous system during development and the fetal onset of motor defect associated with missense variants of *SCN1A* reported here

strongly suggest that *SCN1A* has an essential role in prenatal motor development. Our data show that variants of *SCN1A* encoding essential components of the AIS are responsible for severe developmental motor defects leading to AMC in addition to epilepsy. Taken together, the results of our study combined with previous investigations suggest a large spectrum of *SCN1A*-related phenotypes ranging from early-onset epileptic encephalopathy to fetal onset of motor defect leading to arthrogryposis.

SCN1A variants in Dravet syndrome have been mainly associated with loss-of-function properties. A more severe and earlier onset phenotype has been recently described, associated with p.Thr226Met variant and gain of function properties.²⁶ Importantly, in family 1, both arthrogryposis and early-onset epileptic encephalopathy were observed which suggests that the pathogenic mechanism of the identified missense mutation (p.Leu893Phe) may be more deleterious in our patients leading to severe motor defect in addition to early-onset epilepsy. Similarly, in the patient reported by Spagnoli *et al*,¹⁹ both severe motor defect and epilepsy were observed: this patient carries a de novo *SCN1A* missense mutation (p.Ser228Pro) which reinforces the hypothesis of the same but more severe pathogenic mechanism in this new disease phenotype.

Interestingly, two other genes encoding channels are responsible for similar clinical features. *NALCN* encoding a G-protein-coupled receptor-activated channel is highly expressed in the central nervous system and de novo variants of *NALCN* are responsible for AMC associated with developmental delay (CLIFAHDD, [MIM: 616266]).⁴ Epilepsy was reported in two out of seven affected AMC individuals.⁴ Similarly, *CACNA1E* encodes the alpha1-subunit of the voltage-gated Cav2.3 channel which is highly expressed in the central nervous system. De novo variants of this gene are responsible for early-onset pharmacoresistant seizures (EIEE69, [MIM: 618285]) and 43% of patients had congenital joint contractures.⁵ Altogether, our data enlarge the group of AMC linked to central nervous system involvement caused by variants of genes encoding channels indicating a critical role of these genes in fetal motor development as reported in other developmental channelopathies responsible for cerebral cortex malformations.²⁷

The identification of *SCN1A* pathogenic variants should allow critical information of families on the large clinical spectrum in addition to AMC, such as early-onset epileptic encephalopathy, and the need for careful postnatal follow-up of the affected individuals in order to propose optimal care of both AMC and epilepsy.

Author affiliations

¹Institut National de la Santé et de la Recherche Médicale (Inserm), UMR-1195, Université Paris Saclay, Le Kremlin Bicêtre, 94276, France

²Service de Neurophysiologie Clinique, Centre de référence des maladies neuromusculaires, Hôpital Necker Enfants Malades, Assistance Publique-Hôpitaux de Paris, Université de Paris, Paris, 75015, France

³Service de Génétique, Unité de Génétique Clinique, CHRU de Tours, Hôpital Bretonneau, Tours, 37044, France

⁴Laboratoire de Pathologie, CHU de Rouen and Normandie Univ, Rouen, 76000, France

⁵INSERM, Laboratoire NeoVasc ERI28, Rouen, 76000, France

⁶Neurophysiology Department, Necker-Enfants Malades Hospital, Assistance publique-Hôpitaux de Paris, Paris, 75015, France

⁷Département d'Obstétrique et Gynécologie, Hôpital Antoine-Béclère, Assistance publique-Hôpitaux de Paris, Clamart, 92140, France

⁸Département de Pathologie, CHU Sainte-Justine, Université de Montréal, Québec, H3T 1C5, Québec, Canada

⁹Unité de Foetopathologie, Hôpital Antoine-Béclère, Assistance publique-Hôpitaux de Paris, Clamart, 92140, France

¹⁰Unité de Génétique Médicale, Centre de référence des anomalies du développement et syndromes malformatifs d'Île-de-France, Assistance publique-Hôpitaux de Paris, Le Kremlin Bicêtre, 94276, France

Contributors DJ, DB and MVS conducted molecular analyses. CG, SB, FM, JS, CF-B, JMa and AL recruited and phenotyped the patients. CG conducted electrophysiological studies and contributed to writing the manuscript. AK conducted EEG studies and contributed to writing the manuscript. AL contributed to writing the manuscript. JMe designed the study, performed bioinformatics analysis of whole-exome sequencing data and wrote the manuscript. All authors reviewed the manuscript.

Funding This work was supported by a grant from the Association Française contre les Myopathies (AFM, R16070LL), the Agence de Biomedecine (2016) and the Institut National de la Santé et de la Recherche Médicale (Inserm) to JMe.

Competing interests None declared.

Patient consent for publication Parental/guardian consent obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement All data relevant to the study are included in the article or uploaded as online supplementary information. All data relevant to the study are included in the article and are available.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Judith Melki <http://orcid.org/0000-0002-9125-3171>

REFERENCES

- Hall JG. Genetic aspects of arthrogryposis. *Clin Orthop Relat Res* 1985;44–53.
- Fahy MJ, Hall JG. A retrospective study of pregnancy complications among 828 cases of arthrogryposis. *Genet Couns* 1990;1:3–11.
- Hall JG. Arthrogryposis (multiple congenital contractures): diagnostic approach to etiology, classification, genetics, and general principles. *Eur J Med Genet* 2014;57:464–72.
- Chong JX, McMillin MJ, Shively KM, Beck AE, Marvin CT, Armenteros JR, Buckingham KJ, Nkinsi NT, Boyle EA, Berry MN, Bocian M, Foulds N, Uzielli MLG, Haldeman-Englert C, Hennekam RCM, Kaplan P, Kline AD, Mercer CL, Nowaczyk MJM, Klein Wassink-Ruiter JS, McPherson EW, Moreno RA, Scheuerle AE, Shashi V, Stevens CA, Carey JC, Monteil A, Lory P, Tabor HK, Smith JD, Shendure J, Nickerson DA, Bamshad MJ, University of Washington Center for Mendelian Genomics. De novo mutations in *NALCN* cause a syndrome characterized by congenital contractures of the limbs and face, hypotonia, and developmental delay. *Am J Hum Genet* 2015;96:462–73.
- Helbig KL, Lauerer RJ, Bahr JC, Souza IA, Myers CT, Uysal B, Schwarz N, Gandini MA, Huang S, Keren B, Mignot C, Afejar A, Billette de Villemeur T, Héron D, Nava C, Valence S, Buratti J, Fagerberg CR, Soerensen KP, Kibaek M, Kamsteeg E-J, Koolen DA, Gunning B, Schelhaas HJ, Krueger MC, Fox J, Bakhtiari S, Jarrar R, Padilla-Lopez S, Lindstrom K, Jin SC, Zeng X, Bilguvar K, Papavasiliou A, Xing Q, Zhu C, Boysen K, Vairo F, Lanpher BC, Klee EW, Tillema J-M, Payne ET, Cousin MA, Kruisselbrink TM, Wick MJ, Baker J, Haan E, Smith N, Sadeghpour A, Davis EE, Katsanis N, Corbett MA, MacLennan AH, Gecz J, Biskup S, Goldmann E, Rodan LH, Kichula E, Segal E, Jackson KE, Asamoah A, Dimmock D, McCarrier J, Botto LD, Filloux F, Tvrdik T, Cascino GD, Klingerman S, Neumann C, Wang R, Jacobsen JC, Nolan MA, Snell RG, Lehnert K, Sadleir LG, Anderlid B-M, Kvarnang M, Guerrini R, Friez MJ, Lyons MJ, Leonhard J, Kringlen G, Casas K, El Achkar CM, Smith LA, Rotenberg A, Poduri A, Sanchis-Juan A, Carss KJ, Rankin J, Zeman A, Raymond FL, Blyth M, Kerr B, Ruiz K, Urquhart J, Hughes I, Banka S, Hedrich UBS, Scheffer IE, Helbig I, Zamponi GW, Lerche H, Mefford HC, Task Force for Neonatal Genomics, Deciphering Developmental Disorders Study. De novo pathogenic variants in *CACNA1E* cause developmental and epileptic encephalopathy with contractures, macrocephaly, and dyskinesias. *Am J Hum Genet* 2018;103:666–78.
- Claes L, Del-Favero J, Ceulemans B, Lagae L, Van Broeckhoven C, De Jonghe P. De novo mutations in the sodium-channel gene *SCN1A* cause severe myoclonic epilepsy of infancy. *Am J Hum Genet* 2001;68:1327–32.
- Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* 2009;25:1754–60.
- Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R, 1000 Genome Project Data Processing Subgroup. The sequence alignment/map format and SAMtools. *Bioinformatics* 2009;25:2078–9.
- Wang K, Li M, Hakonarson H. ANNOVAR: functional annotation of genetic variants from high-throughput sequencing data. *Nucleic Acids Res* 2010;38:e164.

- 10 Adzhubei IA, Schmidt S, Peshkin L, Ramensky VE, Gerasimova A, Bork P, Kondrashov AS, Sunyaev SR. A method and server for predicting damaging missense mutations. *Nat Methods* 2010;7:248–9.
- 11 Kumar P, Henikoff S, Ng PC. Predicting the effects of coding non-synonymous variants on protein function using the SIFT algorithm. *Nat Protoc* 2009;4:1073–81.
- 12 Zhou J, Tawk M, Tiziano FD, Veillet J, Bayes M, Nolent F, Garcia V, Servidei S, Bertini E, Castro-Giner F, Renda Y, Carpentier S, Andrieu-Abadie N, Gut I, Levade T, Topaloglu H, Melki J. Spinal muscular atrophy associated with progressive myoclonic epilepsy is caused by mutations in *ASAH1*. *Am J Hum Genet* 2012;91:5–14.
- 13 Miller AR, Hawkins NA, McCollom CE, Kearney JA. Mapping genetic modifiers of survival in a mouse model of Dravet syndrome. *Genes Brain Behav* 2014;13:163–72.
- 14 Nelson AD, Jenkins PM. Axonal membranes and their domains: assembly and function of the axon initial segment and node of Ranvier. *Front Cell Neurosci* 2017;11:136.
- 15 Catterall WA. From ionic currents to molecular mechanisms: the structure and function of voltage-gated sodium channels. *Neuron* 2000;26:13–25.
- 16 Duflocq A, Le Bras B, Bullier E, Couraud F, Davenne M. Nav1.1 is predominantly expressed in nodes of Ranvier and axon initial segments. *Mol Cell Neurosci* 2008;39:180–92.
- 17 Jones SL, Svitkina TM. Axon initial segment cytoskeleton: architecture, development, and role in neuron polarity. *Neural Plast* 2016;2016:1–19.
- 18 Gitiaux C, Chemaly N, Quijano-Roy S, Barnerias C, Desguerre I, Hully M, Chiron C, Dulac O, Nabbout R. Motor neuropathy contributes to crouching in patients with Dravet syndrome. *Neurology* 2016;87:277–81.
- 19 Spagnoli C, Frattini D, Rizzi S, Salerno GG, Fusco C. Early infantile SCN1A epileptic encephalopathy: expanding the genotype–phenotype correlations. *Seizure* 2019;65:62–4.
- 20 Laquerrière A, Maluenda J, Camus A, Fontenas L, Dieterich K, Nolent F, Zhou J, Monnier N, Latour P, Gentil D, Héron D, Desguerre I, Landrieu P, Beneteau C, Delaporte B, Bellesme C, Baumann C, Capri Y, Goldenberg A, Lyonnet S, Bonneau D, Estournet B, Quijano-Roy S, Francannet C, Odent S, Saint-Frison M-H, Sigaudy S, Figarella-Branger D, Gelot A, Mussini J-M, Lacroix C, Drouin-Garraud V, Malinge M-C, Attié-Bitach T, Bessieres B, Bonniere M, Encha-Razavi F, Beaufrère A-M, Khung-Savatovsky S, Perez MJ, Vasiljevic A, Mercier S, Roume J, Trestard L, Saugier-Verber P, Cordier M-P, Layet V, Legendre M, Vigouroux-Castera A, Lunardi J, Bayes M, Jouk PS, Rigonnot L, Granier M, Sternberg D, Warszawski J, Gut I, Gonzales M, Tawk M, Melki J. Mutations in *CNTNAP1* and *ADCY6* are responsible for severe arthrogryposis multiplex congenita with axoglial defects. *Hum Mol Genet* 2014;23:2279–89.
- 21 Maluenda J, Manso C, Quevarec L, Vivanti A, Marguet F, Gonzales M, Guimiot F, Petit F, Toutain A, Whalen S, Grigorescu R, Coeslier AD, Gut M, Gut I, Laquerrière A, Devaux J, Melki J. Mutations in *GLDN*, encoding gliomedin, a critical component of the nodes of Ranvier, are responsible for lethal arthrogryposis. *Am J Hum Genet* 2016;99:928–33.
- 22 Wallace RH, Scheffer IE, Barnett S, Richards M, Dibbens L, Desai RR, Lerman-Sagie T, Lev D, Mazarib A, Brand N, Ben-Zeev B, Goikhman I, Singh R, Kremmidoitis G, Gardner A, Sutherland GR, George AL, Mulley JC, Berkovic SF. Neuronal sodium-channel alpha1-subunit mutations in generalized epilepsy with febrile seizures plus. *Am J Hum Genet* 2001;68:859–65.
- 23 Dichgans M, Freilinger T, Eckstein G, Babini E, Lorenz-Depiereux B, Biskup S, Ferrari MD, Herzog J, van den Maagdenberg AMJM, Pusch M, Strom TM. Mutation in the neuronal voltage-gated sodium channel *SCN1A* in familial hemiplegic migraine. *Lancet* 2005;366:371–7.
- 24 Symonds JD, Zuberi SM, Stewart K, McLellan A, O'Regan M, MacLeod S, Jollands A, Joss S, Kirkpatrick M, Brunklaus A, Pilz DT, Shetty J, Dorris L, Abu-Arafah I, Andrew J, Brink P, Callaghan M, Cruden J, Diver LA, Findlay C, Gardiner S, Grattan R, Lang B, MacDonnell J, McKnight J, Morrison CA, Nairn L, Slean MM, Stephen E, Webb A, Vincent A, Wilson M. Incidence and phenotypes of childhood-onset genetic epilepsies: a prospective population-based national cohort. *Brain* 2019;142:2303–18.
- 25 Catarino CB, Liu JYW, Liagkouras I, Gibbons VS, Labrum RW, Ellis R, Woodward C, Davis MB, Smith SJ, Cross JH, Appleton RE, Yendle SC, McMahon JM, Bellows ST, Jacques TS, Zuberi SM, Koepp MJ, Martinian L, Scheffer IE, Thom M, Sisodiya SM. Dravet syndrome as epileptic encephalopathy: evidence from long-term course and neuropathology. *Brain* 2011;134:2982–3010.
- 26 Berecki G, Bryson A, Terhag J, Maljevic S, Gazina EV, Hill SL, Petrou S. SCN1A gain of function in early infantile encephalopathy. *Ann Neurol* 2019;85:514–25.
- 27 Smith RS, Walsh CA. Ion channel functions in early brain development. *Trends Neurosci* 2020;43:103–14.