OPEN ACCESS

UK recommendations for *SDHA* germline genetic testing and surveillance in clinical practice

Helen Hanson ,^{1,2} Miranda Durkie,³ Fiona Laloo,⁴ Louise Izatt ,⁵ Terri P McVeigh,⁶ Jackie A Cook,⁷ Carole Brewer,⁸ James Drummond,⁹ Samantha Butler,¹⁰ Treena Cranston,¹¹ Ruth Casey,^{12,13} Tricia Tan,¹⁴ Daniel Morganstein,¹⁵ Diana M Eccles ,¹⁶ Marc Tischkowitz,¹³ Clare Turnbull,² Emma Roisin Woodward ,^{4,17} Eamonn R Maher ,¹³ on behalf of UK Cancer Genetics Centres

For numbered affiliations see end of article.

Correspondence to

Dr Helen Hanson, South West Thames Regional Genetics Service, St George's University Hospitals NHS Foundation Trust, London SW17 0QT, UK; helen.hanson@stgeorges.nhs.uk

Received 22 November 2021

Accepted 13 February 2022

ABSTRACT

SDHA pathogenic germline variants (PGVs) are identified in up to 10% of patients with paraganglioma and pheochromocytoma and up to 30% with wild-type gastrointestinal stromal tumours. Most *SDHA* PGV carriers present with an apparently sporadic tumour, but often the pathogenic variant has been inherited from parent who has the variant, but has not developed any clinical features. Studies of *SDHA* PGV carriers suggest that lifetime penetrance for *SDHA*-associated tumours is low, particularly when identified outside the context of a family history. Current recommended surveillance for *SDHA* PGV carriers follows an intensive protocol. With increasing implementation of tumour and germline large panel and whole-genome sequencing, it is likely more *SDHA* PGV carriers will be identified in patients with tumours not strongly associated with *SDHA*, or outside the context of a strong family history. This creates a complex situation about what to recommend in clinical practice considering low penetrance for tumour development, surveillance burden and patient anxiety. An expert *SDHA* working group was formed to discuss and consider this situation. This paper outlines the recommendations from this working group for testing and management of *SDHA* PGV carriers in clinical practice.

INTRODUCTION

SDHA likely pathogenic or pathogenic germline variants (ie, class 4 or class 5 variants according to ACMG/AMP criteria and henceforth collectively referred to as PGVs) are identified in up to 10% of patients with pheochromocytoma and paraganglioma (PPGL) and can account for up to 50% of SDH-deficient wild-type gastrointestinal stromal tumours (wtGISTs) or up to 30% of wtGIST.¹ wtGIST are a unique and uncommon subtype of GISTs that lack somatic-activating variants in the tyrosine kinase c-KIT or platelet-derived growth factor receptor alpha receptors. Most individuals with a PGV in *SDHA* (henceforth referred to as *SDHA* PGV carriers) present with an apparently sporadic tumour without relevant family history, and most PGVs are inherited from a parent who has not presented with any clinical features.²

A large study of patients with paraganglioma (PGL) from the Netherlands, including 30 index

SDHA PGV carriers and 56 non-index *SDHA* PGV carriers, highlighted that the tumour penetrance is low in non-index *SDHA* PGV carriers, with penetrance at age 70 years estimated to be 10%.³ In addition, though penetrance in *SDHA* PGV carriers was estimated as 39% at 40 years of age in a prospective study of a population-based registry of patients with PPGL, there was a significant difference in index patients (45%, n=29) versus *SDHA* carrier relatives (13%, n=9; p<0.001).⁴ Lower estimates of penetrance (~1.7%) have been reported using a Bayesian approach looking at *SDHA* PGVs in EXAC data and patients with PGL and in a cohort of *SDHA* PGV carriers (~0.1%–4.9%).^{5,6} *SDHA* NM_004168.4: c.91C>Tp.(Arg31*) accounts for a large proportion of known carriers and also occurs at frequency of 0.04% (53/128 900) in non-Finnish Europeans (<https://gnomad.broadinstitute.org/variant/5-223624-C-T>, accessed 15 November 2021).³ Despite this frequency there is currently no evidence to suggest that *SDHA* c.91C>Tp.(Arg31*) has a different penetrance to any other *SDHA* PGVs and variant specific management is not advocated by other groups.⁷

While published literature suggests that most *SDHA* PGV carriers will not manifest *SDHA*-related tumours, thus calling into question the clinical utility of surveillance in this low-risk group, probands with *SDHA*-related tumours and confirmed *SDHA* PGVs typically present at young ages (median age at diagnosis is 28 years with a range of 8–76 years⁴). In addition to malignant wtGIST, *SDHA* PGV carriers may also develop malignant PGL.⁸ As for many other rare cancer predisposition genes, there is sparse information available on the effectiveness of surveillance in *SDHA* PGV carriers and the impact of early detection on clinical outcomes.⁹

At the UKCGG (UK Cancer Genetics Group) Consensus meeting in Cambridge in Spring 2019 (<https://www.ukcgg.org/information-education/ukcgg-consensus-meetings/>), a surveillance protocol for *SDHA* PGV carriers was agreed consisting of annual clinical review and biochemistry with abdominal imaging and MRI neck, thorax, abdomen and pelvis at baseline, followed by 3–5 yearly imaging, based on published recommendations and expert opinion.^{10,11}

Since that meeting, questions regarding the utility of predictive testing and surveillance for a

© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY. Published by BMJ.

To cite: Hanson H, Durkie M, Laloo F, et al. *J Med Genet* Epub ahead of print: [please include Day Month Year]. doi:10.1136/jmedgenet-2021-108355

Position statement

Table 1 Succinate-deficient tumours associated with *SDHA* PGV (SDHA-associated/on-target tumours)

Tumour type	Strength of association
Wild-type GIST	+++
Paranglioma*	++
Phaeochromocytoma*	++
Renal cancer†	+
Neuroblastoma‡	Rare
Pituitary adenoma‡	Rare

*For PPGL it is assumed that PGV in other PPGL predisposition genes have been excluded. The National Test Directory now indicates that this testing should include *FH*, *MAX*, *MEN1*, *RET*, *SDHA*, *SDHAF2*, *SDHB*, *SDHC*, *SDHD*, *TMEM127* and *VHL* including analysis for CNVs (National Test Directory indication R223 Inherited phaeochromocytoma and PGL).¹⁷

†For renal cancer, histopathological examination for characteristic features and immunohistochemistry can be helpful in assessing dSDH status.²⁶ SDHB expression is lost in most dSDH tumours with a germline *SDHA*, *SDHB*, *SDHC* or *SDHD* pathogenic variant. Limitation to the utility of SDHB IHC include interobserver variation, false-negative results (presence of SDHB on IHC where a germline variant exists may be more common for *SDHA*-mutated tumours) and equivocal SDHB staining patterns in the presence of germline or somatic *VHL* inactivation.¹¹ SDHA IHC can reveal loss of SDHA expression in SDHA-mutated tumours but is less widely available than SDHB IHC.

‡For neuroblastoma and pituitary adenoma, immunohistochemical evidence of SDH deficiency should be sought and other causes excluded before an *SDHA* variant is considered causal.

GIST, gastrointestinal stromal tumour; IHC, immunohistochemistry; PGL, paraganglioma; PGV, pathogenic germline variant; PPGL, phaeochromocytoma and paraganglioma.

low penetrance condition have been raised in national forums and highlighted by *SDHA* PGVs being identified as secondary findings through the 100,000 genomes project or through wider panel testing in probands with phenotypes not directly related to *SDHA*.¹² It was therefore agreed timely to revisit predictive testing and surveillance guidelines specifically for *SDHA* to address these complex clinical issues.

METHODS

A preliminary scoping survey was sent out to the 24 UK Regional Genetics centres in July 2020, to establish current practice. There was a total of 24 individual responses from 18 centres.

The results of the survey were collated and discussed further by the *SDHA* working group in a virtual meeting and draft recommendations proposed following this meeting. These recommendations were then circulated to three endocrinologists and subsequently to a representative from each of the 24 Regional Genetics centres. Further input was sought from UK laboratories undertaking *SDHA* testing, specifically regarding reporting of *SDHA* PGVs and the final recommendations agreed by the working group.

RECOMMENDATIONS

For these recommendations, it was agreed to define clear 'on-target' *SDHA*-associated tumours based on published literature and expert group opinion to make practical recommendations regarding reporting, clinical management and predictive testing (the terms on-target and SDHA-associated tumours are synonymous, but the term SDHA associated has been used through this document).^{4 13–15} Tumours not specifically listed in this table are currently considered to be 'off-target' tumours (table 1). While it is recognised that *SDHA* PGVs may potentially contribute to a wider phenotypic tumour spectrum, current evidence for further clear associations beyond the tumours defined in table 1 is limited.¹³

Recommendation 1: identification of an *SDHA* PGV in an individual with an *SDHA*-associated tumour

When diagnostic genetic testing is undertaken in an individual with wtGIST, PPGL, renal cancer, neuroblastoma or pituitary tumour and a PGV identified, including *SDHA* c.91C>Tp. (Arg31*), provided other causes have been excluded or there is appropriate immunohistochemical evidence (see table 1 footnotes), the PGV can be considered to be associated with the clinical phenotype and the diagnostic laboratory report should reflect this. However, we would recommend that the report highlights the low penetrance of *SDHA* PGVs and the need for onward referral to a clinical genetics service for the discussion of predictive genetic testing (table 2). Suggested wording approved by the UK Can-VIG group¹⁶ is 'SDHA pathogenic germline variants appear to have a very low penetrance in asymptomatic relatives who are heterozygous for the variant. Therefore, while predictive testing in other family members may be offered,

Table 2 Summary of recommendations

	Report germline likely pathogenic/ pathogenic variant in diagnostic setting	Surveillance* for affected proband	Offer predictive testing	Offer surveillance* if positive predictive test
Individual with <i>SDHA</i> -associated tumour (wtGIST, paraganglioma, phaeochromocytoma), renal cancer, neuroblastoma and pituitary adenoma with immunohistochemical evidence of SDH deficiency, see table 1)	Yes	Yes Offer follow-up for initial tumour and surveillance for metachronous tumours (see recommendation 4)	Recommend for FDR, following detailed discussion	Offer surveillance (see recommendation 6) following detailed discussion regarding current knowledge and limitations
Individual with non- <i>SDHA</i> -associated tumour	Yes, recommend reporting is coupled with a recommendation that SDH IHC is performed and the finding is considered to be a non-actionable secondary finding unless there is immunohistochemical evidence of SDHB/ <i>SDHA</i> loss in the tumour or a family history of <i>SDHA</i> -associated tumours†	Not	Not	N/A

*Surveillance: annual symptom review, blood pressure monitoring, biochemistry with plasma metanephrines and 3–5 yearly imaging of neck, thorax, abdomen and pelvis, preferably with MRI from age 15.

†For individuals with non-*SDHA*-associated tumours identified to have a PGV in *SDHA*, we would consider these a non-clinically actionable finding, unless the tumour is demonstrated to show SDHB/*SDHA* loss or there is a family history of *SDHA*-associated tumours. If either IHC loss or family history is confirmed, then recommendations should shift to that for an individual with an *SDHA*-associated tumour.

FDR, first-degree relative; IHC, immunohistochemical; PGV, pathogenic germline variant; wtGIST, wild-type gastrointestinal stromal tumour.

we would recommend referral to clinical genetics for further discussion'.

Recommendation 2: identification of an *SDHA* PGV in an individual with a non-*SDHA*-associated tumour

In most situations within current UK clinical practice, germline *SDHA* testing will only be requested for an individual with a personal or family history of *SDHA*-associated tumours (currently indications R223 and R363 in the current National Genomic Test Directory, October 2021).¹⁷

However, with increasing use of large germline cancer predisposition gene panels, large somatic solid tumour panels and paired whole-genome sequencing (WGS), *SDHA* PGVs may also be identified in off-tumour settings, that is, in individuals with tumours not listed in table 1.

When an *SDHA* PGV is identified during tumour analysis, the likelihood of it being of germline origin is high in both on-tumour (associated tumour types) and off-tumour (non-associated tumour types) settings. Where an *SDHA* pathogenic variant has been identified in tumour tissue, germline testing has been recommended by the European Society of Medical Oncology Precision Medicine Working Group.¹⁸ However, identification of an *SDHA* PGV in an individual with a non-*SDHA*-associated tumour could be considered a secondary or incidental finding. ACMG Recommendations for Reporting of Incidental Findings in Clinical Exome and Genome Sequencing, which are supported and followed within UK practice, suggest that these findings are not reported.¹⁹

We would suggest that if an *SDHA* PGV is identified in an individual with cancer but in an off-tumour setting, for example, through WGS or extended gene panel testing, that the PGV is reported but coupled with a recommendation that SDH immunohistochemistry (IHC) is performed and the finding considered to be a non-actionable secondary finding unless there is immunohistochemical evidence of SDHB/*SDHA* protein loss or a family history of *SDHA*-associated tumours (table 2).

The working group felt that due to the low penetrance of *SDHA* PGVs outside the context of a personal or family history of *SDHA*-associated tumours, in this situation we would not recommend any surveillance in affected individuals or predictive testing for other family members (table 2). Should there be immunohistochemical evidence of SDHB/*SDHA* loss, or suggestive family history, then we would recommend surveillance and predictive testing be undertaken in line with the recommendations for patients with *SDHA*-associated tumours (see recommendations 4–6 below).

Recommendation 3: Identification of an *SDHA* PGV in an individual without cancer

In some situations where an individual has had germline genetic testing for another indication but does not have a personal history of cancer, an *SDHA* PGV may be identified. ACMG Recommendations for Reporting of Incidental Findings in Clinical Exome and Genome Sequencing, which are supported and followed within UK practice, suggest that these findings are not reported.¹⁹

Recommendation 4: surveillance for *SDHA* PGV carriers affected with *SDHA*-associated tumours

For an *SDHA* PGV carrier with an *SDHA*-associated tumour, with respect to the primary tumour, we would recommend at the very least annual clinical examination to include blood pressure assessment and biochemistry to include plasma metanephrines,

combined with imaging of the original tumour region (eg, abdomen if pheochromocytoma, abdominal PGL, GIST or renal tumour) if complete resection was achieved. Ongoing follow-up, surveillance and discharge for their original diagnosis should be determined by multidisciplinary team decision depending on the clinical details and consideration of relevant published guidelines for the follow-up of PPGL.²⁰

There are minimal data on the occurrence of a second tumour in individuals with an *SDHA* PGV. However, metachronous tumours have been reported, in single case reports and in 4 out of 21 index cases in a study from the Netherlands.^{3,8}

With regards to ongoing surveillance for metachronous tumours, based on UKCGG Consensus guidelines,²¹ we would recommend annual symptom review, blood pressure monitoring and biochemistry with plasma metanephrines (with 24-hour urinary metanephrines an alternative especially in children). Imaging should include neck, thorax, abdomen and pelvis at baseline, followed by 3–5 yearly surveillance, preferably with MRI, recognising that follow-up of the original tumour region may be more frequent. Ongoing follow-up should take place in an Endocrinology clinic or Joint Endocrine-Genetics clinic.

The UKCGG guidelines did not make recommendations for pituitary surveillance. This is in part due to the low penetrance for pituitary adenoma, but also the relatively high rate of incidental detection of non-functional microadenoma, estimated to be 10% or greater in a healthy population.²² At present, we would not recommend routine pituitary imaging, in line with a recent international consensus guideline on surveillance in SDHx PGV carriers.⁷ Consideration of annual prolactin and IGF1 and requesting that the MRI neck includes a cut through the pituitary has been proposed as a surveillance recommendation.¹¹ At present, we would only recommend this within the context of a trial or service evaluation and suggest that recommendations are revisited in the future as more data become available.

Recommendation 5: predictive *SDHA* testing

A limited number of small studies have demonstrated that the penetrance of *SDHA* in non-probands is likely to be low: 10%–13% lifetime risk in two studies.^{3,4} These data have raised the question of the utility of predictive testing and surveillance for this patient group. Concern has been raised about the potential to increase anxiety for these families when lifetime risk of an *SDHA*-associated tumour may be low. However, patients can present with aggressive and metastatic disease which may be avoided by early detection. The lifetime penetrance in non-proband family members is also higher than in penetrance estimates from comparing *SDHA* allele frequencies in affected individuals and population controls, ~2%.^{5,6}

The survey of UK Geneticists and Endocrinologists suggested that at present while there is recognition that the cancer risk for non-proband family members may be low, it was felt that the data in this area are limited and there was concern about not discussing predictive testing with family members. There was agreement that while there are limited data available, predictive testing and surveillance should be discussed within the context of our current knowledge, but that this should be reviewed as and when new data on penetrance and effectiveness of surveillance are available.

We would recommend that predictive genetic testing should be considered in families where an *SDHA* PGV has been identified in an individual with an *SDHA*-associated tumour (see tables 1 and 2) or an individual with an off-target tumour not typically associated with *SDHA*, but with suggestive IHC loss

or family history of SDHA-associated tumours (table 2). Due to the reported low penetrance and few reports of familial cases we suggest consideration of offering predictive testing only to first-degree relatives of an affected proband, unless there is a wider family history of SDHA-associated tumours (table 2).

We suggest that detailed discussion regarding predictive testing with these families should take place, ensuring patients are actively involved in the decision-making process regarding predictive testing and a joint decision made whether to proceed. We would recommend particular attention and clear discussion of the low penetrance of SDHA PGVs, counselling of the lack of clear evidence on utility of surveillance and the potential for incidental findings.

We would also recommend detailed discussion of symptoms relating to SDHA-associated tumours, both as an adjunct or alternative to surveillance. We suggest not actively offering predictive testing to second-degree relatives; however, we would recommend providing information on symptoms relating to SDHA-associated tumours and the importance of seeking a specialist opinion if they have specific symptoms, for example, hypertension, severe recurrent headaches, dyspepsia or upper gastrointestinal bleeding or unintentional weight loss.

Recommendation 6: surveillance for unaffected SDHA PGV carriers

As for many other cancer predisposition syndromes, comprehensive data regarding the clinical utility of surveillance are limited in SDHA PGV carriers. However, it is recognised that patients can present at a young age with metastatic disease and cases of positive surveillance have been reported.^{9,23} There is controversy both over whether surveillance should be offered and if it is, the extent of the surveillance. Surveillance with plasma metanephrines alone may not detect the non-secretory head and neck PGL and therefore, if surveillance is recommended it should comprise both biochemical studies and imaging including neck, thorax, abdomen and pelvis.

There are no studies assessing the optimal surveillance interval, but an early study found tumour doubling time of 4.2 years for head and neck PGLs²⁴ and 5.8 years in a series of SDHB and SDHD patients with head and neck PGLs, whereas abdominal and thoracic PGLs grew more slowly, doubling at 6.94 and 11.8 years, respectively.²⁵

Based on expert opinion, UKCGG guidelines in 2019 recommended:

- ▶ Annual symptom review, clinical examination to include blood pressure and annual biochemical surveillance with plasma metanephrines (with 24-hour urinary metanephrines an alternative, especially in children) from age 10.
- ▶ Radiological surveillance every 3–5 years of neck, thorax, abdomen and pelvis, preferably with MRI from age 15. Where possible, imaging is best performed in centres with experience of surveillance for PPGL. Ultrasound is not recommended.
- ▶ Routine pituitary imaging is not recommended (see also Recommendation 4: surveillance for SDHA PGV carriers affected with SDHA-associated tumours section).

These recommendations are in line with recently published international consensus guidelines for SDHx PGV carriers, who also advise that by the age of 70 years, if individuals are well with no tumours, the interval of imaging can be increased to 5 yearly until age 80 and then stopped if well.⁷ We would suggest that predictive genetic testing is considered from the age that surveillance commences, that is, around 10 years.

CONCLUSION AND FUTURE WORK

It was recognised that there is currently limited knowledge regarding the full phenotype and penetrance of tumours in individuals with SDHA PGVs. Therefore, there is a critical need for systematic prospective data collection to address this and the outcomes of surveillance.

It was also recognised that there has been inconsistency in practice across the country, and it is hoped that these recommendations will help to address this. To improve patient understanding, development of a national patient leaflet is planned to both highlight the uncertainty regarding true cancer risk for this patient group, the pros and cons of predictive testing and surveillance and education on symptoms.

Given the uncertainties and lack of data highlighted in this report, the working group recommend that:

1. Prospective research studies and/or service evaluations are undertaken to define of the natural history of individuals with SDHA PGVs identified with SDHA-associated and SDHA non-associated tumours and consider if specific genetic or environmental factors alter the penetrance in SDHA carriers.
2. The outcomes and clinical utility of surveillance in SDHA PGV carrier probands and non-probands are carefully documented and evaluated.
3. Novel means of early detection are sought to reduce burden on radiology departments.
4. The opinions and preferences of patient groups are canvassed.

Author affiliations

¹South West Thames Regional Genetic Services, St George's University Hospitals NHS Foundation Trust, London, UK

²Division of Genetics and Epidemiology, Institute of Cancer Research, London, UK

³Sheffield Diagnostic Genetics Service, Sheffield Children's NHS Foundation Trust, North East and Yorkshire Genomic Laboratory Hub, Sheffield, UK

⁴Clinical Genetics Service, Manchester Centre for Genomic Medicine, Central Manchester University Hospitals NHS Foundation Trust, Manchester, UK

⁵Clinical Genetics, Guy's and St Thomas' NHS Foundation Trust, London, UK

⁶Cancer Genetics Unit, Royal Marsden NHS Foundation Trust, London, UK

⁷Department of Clinical Genetics, Sheffield Children's NHS Foundation Trust, Sheffield, UK

⁸Department of Clinical Genetics, Royal Devon and Exeter NHS Foundation Trust, Exeter, UK

⁹East NHS Genomic Laboratory Hub, Cambridge University Hospitals Genomic Laboratory, Cambridge University Hospital Foundation Trust, Cambridge, UK

¹⁰Molecular Genetics, West Midlands Regional Genetics Laboratory, Birmingham, UK

¹¹Oxford Molecular Genetics Laboratory, Churchill Hospital, Oxford, UK

¹²Department of Endocrinology, Cambridge University Hospitals NHS Foundation Trust, Cambridge, UK

¹³Department of Medical Genetics, University of Cambridge and Cambridge University Hospitals NHS Foundation Trust, Cambridge, UK

¹⁴Section of Investigative Medicine, Imperial College London, London, UK

¹⁵Endocrinology, Royal Marsden Hospital NHS Trust, London, UK

¹⁶Cancer Sciences Unit, Faculty of Medicine, University of Southampton, Southampton, UK

¹⁷Division of Evolution and Genomic Sciences, School of Biological Sciences, Faculty of Biology Medicine and Health, The University of Manchester, Manchester, UK

Twitter Clare Turnbull @clare_turnbull and Emma Roisin Woodward @ER_Woodward

Collaborators Representatives from UK Cancer Genetics Centres: Alan Donaldson, Clinical Genetics, Bristol; Frauke Pelz, Clinical Genetics, Cardiff; Angela George, The Royal Marsden NHS Foundation Trust; Anju Kulkarni, Clinical Genetics, Guys and St Thomas' NHS Foundation Trust; Dorothy Halliday, Clinical Genetics, Oxford; Angela Brady, Clinical Genetics, North West Thames Regional Genetics Service; Katie Snape, South-West Thames Regional Genetics Service; Kai-Ren Ong Genetics, Birmingham; Rachel Harrison, Nottingham Clinical Genetics Service; Julian Adlard, Yorkshire Regional Genetics Service, Leeds Teaching Hospitals NHS Trust; Lynn Greenhalgh, Clinical Genetics, Liverpool; Jennie Murray, Edinburgh Genetics Service; Ajith Kumar Clinical Genetics, Great Ormond Street Hospital; Richard Martin Northern Genetics

Service, Newcastle-upon-Tyne; David Goudie Clinical Genetics, Dundee; Rosemarie Davison Clinical Genetics, Glasgow; Gillian Rea Clinical Genetics, Belfast; Zosia Miedzybrodzka, Clinical Genetics, Julian Barwell, Leicester Clinical Genetics Service; Lucy Side Clinical Genetics, Southampton.

Contributors Planning and working group: HH, MD, FL, LI, TPM, JAC, CB, ERW, ERM; Expert input to working group: JD, SB, TC, RC, TT, DM; Manuscript draft: HH; Editing and review of manuscript: HH, MD, FL, LI, TPM, JAC, CB, JD, SB, TC, RC, TT, DM, DME, MT, CT, ERW, ERM; Funding: CT, DME, MT; Scoping survey completion and manuscript review: All named collaborators at UK Cancer Genetics Centres.

Funding HH and FL are supported by Cancer Research CRUK Catalyst Award, CanGene-CanVar (C61296/A27223). RC is supported by GIST Support UK. ERW is supported by the Manchester NIHR Biomedical Research Centre (IS-BRC-1215-20007). ERM thanks the NIHR Cambridge Biomedical Research Centre for support. The University of Cambridge has received salary support (ERM) from the NHS in the East of England through the Clinical Academic Reserve.

Competing interests RC has received speakers' honoraria from Novartis and Ipsen; DM has received speakers' honoraria from Roche and MSD and speakers' honoraria and consulting fees from Bristol Myers Squibb.

Patient consent for publication Not required.

Ethics approval This study does not involve human participants.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

ORCID iDs

Helen Hanson <http://orcid.org/0000-0002-3303-8713>

Louise Izatt <http://orcid.org/0000-0003-1258-4843>

Diana M Eccles <http://orcid.org/0000-0002-9935-3169>

Emma Roisin Woodward <http://orcid.org/0000-0002-6297-2855>

Eamonn R Maher <http://orcid.org/0000-0002-6226-6918>

REFERENCES

- Boikos SA, Pappo AS, Killian JK, LaQuaglia MP, Weldon CB, George S, Trent JC, von Mehren M, Wright JA, Schiffman JD, Raygada M, Pacak K, Meltzer PS, Miettinen MM, Stratakis C, Janeway KA, Helman LJ. Molecular subtypes of KIT/PDGFRA wild-type gastrointestinal stromal tumors. *JAMA Oncol* 2016;2.
- Casey RT, Ascher DB, Rattenberry E, Izatt L, Andrews KA, Simpson HL, Challis B, Park S-M, Bulusu VR, Lalloo F, Pires DEV, West H, Clark GR, Smith PS, Whitworth J, Papatthomas TG, Taniere P, Savisaar R, Hurst LD, Woodward ER, Maher ER. SDHA related tumorigenesis: a new case series and literature review for variant interpretation and pathogenicity. *Mol Genet Genomic Med* 2017;5:237–50.
- van der Tuin K, Mensenkamp AR, Tops CMJ, Corssmit EPM, Dinjens WN, van de Horst-Schrivers ANA, Jansen JC, de Jong MM, Kunst HPM, Kusters B, Leter EM, Morreau H, van Nesselrooij BMP, Oldenburg RA, Spruijt L, Hes FJ, Timmers HJLM. Clinical aspects of SDHA-related pheochromocytoma and paraganglioma: a nationwide study. *J Clin Endocrinol Metab* 2018;103:438–45.
- Bausch B, Schiavi F, Ni Y, Welander J, Patocs A, Ngeow J, Wellner U, Malinoc A, Taschin E, Barbon G, Lanza V, Söderkvist P, Stenman A, Larsson C, Svahn F, Chen J-L, Marquard J, Fraenkel M, Walter MA, Peczkowska M, Prejbisz A, Jarzab B, Hasse-Lazar K, Petersenn S, Moeller LC, Meyer A, Reisch N, Trupka A, Brase C, Galiano M, Preuss SF, Kwok P, Lendvai N, Berisha G, Makay Özer, Boedeker CC, Weryha G, Racz K, Januszewicz A, Walz MK, Gimm O, Opocher G, Eng C, Neumann HPH, European-American-Asian Pheochromocytoma-Paraganglioma Registry Study Group. Clinical characterization of the pheochromocytoma and paraganglioma susceptibility genes SDHA, TMEM127, Max, and SDHAF2 for gene-informed prevention. *JAMA Oncol* 2017;3:1204–12.
- Benn DE, Zhu Y, Andrews KA, Wilding M, Duncan EL, Dwight T, Tothill RW, Burgess J, Crook A, Gill AJ, Hicks R, Kim E, Luxford C, Marfan H, Richardson AL, Robinson B, Schlosberg A, Susman R, Tacon L, Trainer A, Tucker K, Maher ER, Field M, Clifton-Bligh RJ. Bayesian approach to determining penetrance of pathogenic SDH variants. *J Med Genet* 2018;55:729–34.
- Maniamp P, Zhou K, Loneragan M, Berg JN, Goudie DR, Newey PJ. Pathogenicity and Penetrance of Germline SDHA Variants in Pheochromocytoma and Paraganglioma (PPGL). *J Endocr Soc* 2018;2:806–16.
- Amar L, Pacak K, Steichen O, Akker SA, Aylwin SJB, Baudin E, Buffet A, Burnichon N, Clifton-Bligh RJ, Dahia PLM, Fassnacht M, Grossman AB, Herman P, Hicks RJ, Januszewicz A, Jimenez C, Kunst HPM, Lewis D, Mannelli M, Naruse M, Robledo M, Taieb D, Taylor DR, Timmers HJLM, Treglia G, Tufton N, Young WF, Lenders JWM, Gimenez-Roqueplo A-P, Lussey-Lepoutre C. International consensus on initial screening and follow-up of asymptomatic SDHx mutation carriers. *Nat Rev Endocrinol* 2021;17:435–44.
- Tufton N, Ghelani R, Srirangalingam U, Kumar AV, Drake WM, Iacovazzo D, Skordilis K, Berney D, Al-Mrayat Ma'en, Khoo B, Akker SA. SDHA mutated paragangliomas may be at high risk of metastasis. *Endocr Relat Cancer* 2017;24:L43–9.
- White G, Tufton N, Akker SA. First-positive surveillance screening in an asymptomatic SDhA germline mutation carrier. *Endocrinol Diabetes Metab Case Reports* 2019;2019.
- Wong MY, Andrews KA, Challis BG, Park S-M, Acerini CL, Maher ER, Casey RT. Clinical practice guidance: surveillance for pheochromocytoma and paraganglioma in paediatric succinate dehydrogenase gene mutation carriers. *Clin Endocrinol* 2019;90:499–505.
- MacFarlane J, Seong KC, Bisambar C, Madhu B, Allinson K, Marker A, Warren A, Park S-M, Giger O, Challis BG, Maher ER, Casey RT. A review of the tumour spectrum of germline succinate dehydrogenase gene mutations: beyond pheochromocytoma and paraganglioma. *Clin Endocrinol* 2020;93:528–38.
- Genomics England. 100000 genome Project. Available: <https://www.genomicsengland.co.uk/about-genomics-england/the-100000-genomes-project>
- Dubard Gault M, Mandelker D, DeLair D, Stewart CR, Kemel Y, Sheehan MR, Siegel B, Kennedy J, Marcell V, Arnold A, Al-Ahmadie H, Modak S, Robson M, Shukla N, Roberts S, Vijai J, Topka S, Kentsis A, Cadoo K, Carlo M, Latham Schwark A, Reznik E, Dinatale R, Hechtman J, Borrás Flores E, Jairam S, Yang C, Li Y, Bayraktar EC, Ceyhan-Birsoy O, Zhang L, Kohlman W, Schiffman J, Stadler Z, Birsoy K, Kung A, Offit K, Walsh MF. Germline SDHA mutations in children and adults with cancer. *Cold Spring Harb Mol Case Stud* 2018;4:1–12.
- Korpershoek E, Favier J, Gaal J, Burnichon N, van Gessel B, Oudijk L, Badoual C, Gadessaud N, Venisse A, Bayley J-P, van Dooren MF, de Herder WW, Tissier F, Plouin P-F, van Nederveen FH, Dinjens WNM, Gimenez-Roqueplo A-P, de Krijger R. SDhA immunohistochemistry detects germline SDhA gene mutations in apparently sporadic paragangliomas and pheochromocytomas. *J Clin Endocrinol Metab* 2011;96:E1472–6.
- Burnichon N, Brière J-J, Libé R, Vescovo L, Rivière J, Tissier F, Jouanno E, Jeunemaitre X, Bénit P, Tzagoloff A, Rustin P, Bertherat J, Favier J, Gimenez-Roqueplo A-P. SDhA is a tumor suppressor gene causing paraganglioma. *Hum Mol Genet* 2010;19:3011–20.
- Garrett A, Callaway A, Durkie M, Cubuk C, Alikian M, Burghel GJ, Robinson R, Izatt L, Talukdar S, Side L, Cranston T, Palmer-Smith S, Baralle D, Berry IR, Drummond J, Wallace AJ, Norbury G, Eccles DM, Ellard S, Lalloo F, Evans DG, Woodward E, Tischkowitz M, Hanson H, Turnbull C, CanVIG-UK. Cancer variant interpretation group UK (CanVIG-UK): an exemplar national subspecialty multidisciplinary network. *J Med Genet* 2020;57:829–34.
- NHS. National genomic test directory testing criteria for rare and inherited disease. Available: <https://www.england.nhs.uk/wp-content/uploads/2018/08/Rare-and-inherited-disease-eligibility-criteria-2021-22-v2.pdf>
- Mandelker D, Donoghue M, Talukdar S, Bandlamudi C, Srinivasan P, Vivek M, Jezdic S, Hanson H, Snape K, Kulkarni A, Hawkes L, Douillard J-Y, Wallace SE, Rial-Sebbag E, Meric-Bersntam F, George A, Chubb D, Loveday C, Ladanyi M, Berger MF, Taylor BS, Turnbull C. Germline-focussed analysis of tumour-only sequencing: recommendations from the ESMO precision medicine Working group. *Ann Oncol* 2019;30:1221–31.
- Kalia SS, Adelman K, Bale SJ, Chung WK, Eng C, Evans JP, Herman GE, Hufnagel SB, Klein TE, Korf BR, McKelvey KD, Ormond KE, Richards CS, Vlangos CN, Watson M, Martin CL, Miller DT. Recommendations for reporting of secondary findings in clinical exome and genome sequencing, 2016 update (ACMG SF v2.0): a policy statement of the American College of medical genetics and genomics. *Genet Med* 2017;19:249–55.
- Plouin PF, Amar L, Dekkers OM, Fassnacht M, Gimenez-Roqueplo AP, Lenders JWM, Lussey-Lepoutre C, Steichen O, Guideline Working Group. European Society of endocrinology clinical practice guideline for long-term follow-up of patients operated on for a pheochromocytoma or a paraganglioma. *Eur J Endocrinol* 2016;174:G1–10.
- UK CGG. UKCGG consensus meetings. Available: <https://www.ukcgg.org/information-education/ukcgg-consensus-meetings>
- Ntali G, Wass JA, Epidemiology WJA. Epidemiology, clinical presentation and diagnosis of non-functioning pituitary adenomas. *Pituitary* 2018;21:111–8.
- Greenberg SE, Jacobs MF, Wachtel H, Anson A, Buchmann L, Cohen DL, Bonanni M, Bennett B, Naumer A, Schaefer AM, Kohlmann W, Nathanson KL, Else T, Fishbein L. Tumor detection rates in screening of individuals with SDHx-related hereditary paraganglioma-pheochromocytoma syndrome. *Genet Med* 2020;22:1–7.
- Jansen JC, van den Berg R, Kuiper A, van der Mey AG, Zwiderman AH, Cornelisse CJ. Estimation of growth rate in patients with head and neck paragangliomas influences the treatment proposal. *Cancer* 2000;88:2811–6.
- Michałowska I, Cwikła JB, Michalski W, Wyrwicz LS, Prejbisz A, Szperl M, Nieć D, Neumann HPH, Januszewicz A, Peczkowska M. Growth rate of paragangliomas related to germline mutations of the SDHx genes. *Endocr Pract* 2017;23:342–52.
- Kuroda N, Yorita K, Nagasaki M, Harada Y, Ohe C, Jeruc J, Raspolini MR, Michal M, Hes O, Amin MB. Review of succinate dehydrogenase-deficient renal cell carcinoma with focus on clinical and pathobiological aspects. *Pol J Pathol* 2016;67:3–7.