

## **SUPPLEMENTARY MATERIALS AND METHODS**

### **Immunofluorescence staining**

Cells were fixed in 4% paraformaldehyde in PBS for 20 min at room temperature followed by permeabilization in 0.1% Triton X-100 for 5 min. After blocking in 3% BSA (Cat.# 05482, Sigma), cells were incubated in anti-EEA1 (Cat.# 610456, BD) primary antibody for 1 hr at room temperature, followed by incubation for 30 min at room temperature in secondary antibodies (Alexa Fluor® 568, Invitrogen).

### **Generation of knock-out (KO) or knock-down (KD) cell lines**

3T3 and H4 cell lines lacking VPS35L were generated using CRISPR/Cas9 technology. The target sequence was TTTCAAGATCGCTTCCATCCGGG in exon 10 for 3T3, and CAGCGTCTTACCAGGTAATGCGG in exon9-intron9 for H4. Clones were isolated and gene disruption was validated by PCR-based sequencing and western blot. A 3T3 cell line of VPS35L knock-down was generated using VPS35L-shRNA-expressing lentivirus (Merck). 3T3 cells were incubated with virus-containing medium for 24 hr, then cells were cultured in puromycin containing medium for more than two weeks to select for shRNA expressing cells.

### **Statistical analysis**

Results are presented as the mean  $\pm$  standard error measurement (s.e.m). A two-sided Student's

*t*-test was performed to compare the means between two groups. When the means of three groups were to be compared, one-way analysis of variance (ANOVA) with post hoc Tukey's or Dunnet's honestly significant difference calculator test was used. Statistics were calculated using EZR (Saitama Medical Center, Jichi Medical University).  $P < 0.05$  was considered significant.