

Table 1: Sex distribution, age at onset, attack and headache symptoms

	FHM (n=112)	Familial MA (n=487)
Sex distribution (men:women)	1:3.3 (26 M, 86 W)	1:2.3 (148 M, 339 W)
Mean age at onset (years;range)	16.5 (1-42)	21.4 (4-77)
Lifetime number of attacks:		
1	0 (0%)	5 (1%)
2 to 4	18 (16%)	37 (8%)
5 to 9	7 (6%)	37 (8%)
10 to 49	21 (19%)	155 (32%)
50 to 100	19 (17%)	115 (24%)
> 100	46 (41%)	129 (27%)
Duration of attack:		
<30 minuts	0 (0%)	2 (0.5%)
30 min - 4 hours	8 (7%)	119 (26%)
4 h - 24 h	70 (63%)	254 (56%)
24 h - 72 h	32 (29%)	70 (15%)
3 days - 7 days	1 (1%)	8 (2%)
> 7 days	0 (0%)	2 (0.5%)
Headache phase:		
Unilateral	59%	59%
Pulsating	86%	55%
Moderate/severe intensity	99%	85%
Aggravation by physical exercise	97%	80%
Nausea	84%	70%
Vomiting	61%	43%
Photophobia	89%	79%
Phonophobia	87%	58%

FHM: familial hemiplegic migraine. MA: Migraine with typical aura. MO: Migraine without aura.

toms for FHM, familial MA/MO and sporadic MA/MO

Familial MO (n=253)	Sporadic MA (n=750)	Sporadic MO (n=954)
1:1.9 (86 M, 167 W)	1:3.7 (160 M, 587 W)	1:6.2 (133 M, 817 W)
19.4 (0-64)	22.8 (0-70)	21 (2-60)
6 (2%)	5 (1%)	0 (0%)
7 (3%)	41 (6%)	1 (0%)
10 (4%)	39 (5%)	2 (0%)
61 (25%)	161 (22%)	32 (3%)
46 (19%)	108 (15%)	80 (8%)
118 (48%)	385 (52%)	833 (88%)
0 (0%)	6 (1%)	1 (0%)
17 (7%)	73 (10%)	11 (1%)
150 (60%)	292 (42%)	306 (32%)
66 (27%)	271 (39%)	524 (55%)
14 (56%)	56 (8%)	100 (11%)
2 (1%)	3 (1%)	7 (1%)
58%	73%	83%
69%	65%	72%
99%	93%	99%
88%	84%	92%
81%	81%	88%
44%	55%	60%
79%	89%	91%
72%	78%	85%

aura.